

| | Year | Rate | Indicator |
|-------------|--------------|--------------------------------|--------------------------------------------------------------------|
| MOPH, 2011 | PAPFAM, 2004 | 93% | Pregnant women attended by trained personnel |
| | 19 | per 500 000 live births | Facilities with functioning essential obstetric care |
| | PAPFAM, 2004 | 98% | Deliveries attended by trained personnel |
| | MICS3, 2009 | 96% | Deliveries undertaken in health facilities** |
| | MNMNS, 2013 | 17 | Proportion of caesarean sections of all deliveries*** |
| MNMNS, 2013 | MICS3, 2009 | 54% | Married women (15-49) using contraceptives |
| | | | Pregnant women with anaemia |
| | PAPFAM, 2004 | 1.9 | Total fertility rate |
| | 17 | 16.3 | Maternal mortality ratio (maternal deaths per 100 000 live births) |
| | MNMNS, 2013 | 16.3 | Perinatal mortality rate |
| | MNMNS, 2013 | 7.7% | Neonatal mortality rate (number per 1000 live births) |
| MNMNS, 2013 | 7.7% | Percentage of low birth weight | |

Notes:

*Calculated as number of hospitals with required equipments (n=150, population=4 036 450)

**Taking percentage of babies weighed upon birth as quasi indicator

***Maternal neonatal mortality notification system, MOPH

Calculated as (IUCD + stillbirths + deaths in the first week of life), per 1000 births

Monday 6th of May 2024 05:32:16 PM