	☐ Year	Rate	Indicator
	PAPFAM, 2004	9 .P iieg	nant women attended by trained personnel \Box
МОРН, 2011	19	per 500 00 0\pers fo fac i	lities with functioning essential obstetric care
			-
	PAPFAM, 2004	98%	Deliveries attended by trained personnel
	MICS3, 2009		Deliveries undertaken in health facilities**
	MNMNS, 2013		on of caesarean sections of all deliveries***
	MICS3, 2009	54% N	Married women (15-49) using contraceptives☐
			Pregnant women with anaemia
	PAPFAM, 2004	1.9	Total fertility rate☐
MNMNS, 2013	17	Materbiaths)	tality ratio (maternal deaths per 100 000 live
			ъ П
	MNMNS, 2013	16.3	Perinatal mortality rate
	MNMNS, 2013		mortality rate (number per 1000 live births) Percentage of low birth weight
	MNMNS, 2013	7.7%	Percentage of low birth weight⊔
			Notes:
		*Calculated as number of hospitals with req	uired equipments (n=150, population=4 036 450)
		**Taking percentage	e of babies weighed upon birth as quasi indicator
		***Matern:	al neonatal mortality notification system, MOPH

Calculated as (IUFD + stillbirths + deaths in the first week of life), per 1000 births

Sunday 27th of April 2025 07:58:33 PM