


TIME TO VACCINATE

If you are offered one of these COVID-19 vaccines, know when you should take the second dose.

Pfizer	two doses (0.3 ml each) given intramuscularly into the deltoid muscle. An interval of 21-28 days between the doses is recommended.
Moderna	two doses (0.5 ml each) given intramuscularly into the deltoid muscle. An interval of 28 days between doses is recommended.
AstraZeneca	two doses (0.5 ml) given intramuscularly into the deltoid muscle. An interval of 8 to 12 weeks between the doses is recommended.
Janssen	one dose (0.5 ml) given intramuscularly into the deltoid muscle.
Sinopharm	two doses (0.5 ml) given intramuscularly into the deltoid muscle. An interval of 3-4 weeks between the doses.



World Health Organization
www.who.int/emro

[Download](#)

WHAT YOU NEED TO KNOW ABOUT THE ASTRAZENECA VACCINE

OVER 200 MILLION DOSES ADMINISTERED WORLDWIDE*

VERY EFFECTIVE IN PREVENTING SEVERE COVID-19 INFECTION, LONG-TERM COMPLICATIONS AND DEATH

PREVENTS THE RISK OF SEVERE DISEASE FROM SOME VARIANT STRAINS OF THE VIRUS

A VERY RARE SIDE EFFECT (BLOOD CLOTTING WITH LOW PLATELETS) MAY OCCUR AFTER A FIRST DOSE

THESE VERY RARE SIDE EFFECTS OCCUR IN 4-6 PEOPLE IN A MILLION AND ARE LESS LIKELY IN OLDER ADULTS


THE RISK OF BLOOD CLOTS DUE TO COVID-19 IS MUCH HIGHER THAN RISK OF CLOTS DUE TO VACCINE

[Download](#) |

[French](#)

The truth about COVID-19 vaccines


COVID-19 vaccines trigger the body's natural immunity to respond to the virus, so that it can fight real infection in the future.




[Download](#)




[Download](#)



[Download](#)



[Download](#)



[Download](#)



[Download](#)



[Download](#)



[Download](#)

Friday 17th of May 2024 10:38:45 AM