18 January 2023, Cairo, Egypt – The recently detected XBB.1.5 variant is a sub-lineage of the Omicron XBB sub-lineage. Although data on XBB.1.5 are still very limited, currently available information indicates that XBB.1.5 has a growth advantage compared to other circulating Omicron sublineages and may therefore contribute to an increase in cases globally.

Additionally, as per preliminary laboratory-based studies, this sublineage evades antibodies in humans at equal rate as XBB* variants are the most antibody-resistant variants to date. No data are currently available regarding the severity of disease associated with XXB.1.5 but assessments are ongoing.

Between 22 October 2022 and 11 January 2023, 5288 sequences of XBB.1.5 have been reported from 38 countries. The majority of these sequences are from the United States of America (82.2%), United Kingdom of Great Britain and Northern Ireland (8.1%), and Denmark (2.2%). Our understanding of the transmissibility, immune escape and severity of disease will increase as more data become available from additional countries. To date, 2 countries in the Eastern Mediterranean Region have reported Omicron subvariant XBB.1.5

WHO's Technical Advisory Group on Virus Evolution (TAG-VE) met on 5 January 2023 to discuss the latest evidence on XBB.1.5 and review the public health risks associated with this variant. As part of its recommendations, TAG-VE urged Member States to expand sequencing and reinforce surveillance to detect and track the emergence and spread of SARS-CoV-2 new variants, such as XBB.1.5. This will significantly allow well-informed conclusions about their transmissibility, immune escape, severity, and impact on response interventions such as diagnostics, therapeutics and vaccines.

Updates in WHO mask use recommendations

On 13 January 2023, WHO updated its guidelines on mask wearing in community settings, COVID-19 treatments, and clinical management. WHO continues to recommend the use of masks by the public in specific situations, regardless of the local epidemiological situation, given the current spread of the COVID-19 globally. Previously, WHO recommendations were based on the epidemiological situation. Masks are recommended following recent exposure to COVID-19, when someone has or suspects they have COVID-19, when someone is at high-risk of severe COVID-19, and for anyone in a crowded, enclosed, or poorly ventilated space.

Updates in isolation period for COVID-19 patients [

For patients with symptoms, the new guidelines suggest 10 days of isolation from the date of symptom onset. Previously, WHO advised that patients be discharged 10 days after symptom onset, plus at least 3 additional days since their symptoms had resolved.

For those who test positive for COVID-19 but do not have any signs or symptoms, WHO now suggests 5 days of isolation, compared to 10 days previously.

The patient can be discharged from isolation early if they test negative on an antigen-based rapid test.

Isolation of people with COVID-19 is an important step in preventing others from being infected. This can be done at home or at a dedicated facility, such as a hospital or clinic.

The evidence considered by the guideline development group showed that people without symptoms are much less likely to transmit the virus than those with symptoms. Although of very low certainty, evidence also showed that people discharged at day 5 following symptom onset risked infecting 3 times more people than those discharged at day 10.

Review of COVID-19 treatments

WHO has extended its strong recommendation for the use of nirmatrelvir-ritonavir (also known by its brand name 'Paxlovid'). Nirmatrelvir-ritonavir was first recommended by WHO in April 2022. WHO strongly recommends its use in mild or moderate COVID-19 patients who are at high risk of hospitalization.

Pregnant or breastfeeding women with non-severe COVID-19 should consult with their doctor to determine whether they should take this drug, due to 'likely benefits' and a lack of adverse events having been reported.

WHO also reviewed the evidence on 2 other medicines, sotrovimab and casirivimab-imdevimab, and maintains strong recommendations against their use for treating COVID-19. These

monoclonal antibody medicines lack or have diminished activity against the current circulating virus variants.

There are currently 6 proven treatment options for patients with COVID-19, 3 that prevent hospitalization in high-risk persons and 3 that save lives in those with severe or critical disease. Except for corticosteroids, access to other drugs remains unsatisfactory globally.

Friday 25th of April 2025 10:16:21 AM