

### Pin-up The Smoker's Body

Every 15 seconds someone dies from tobacco use, largely from preventable lung cancer. The World Health Organization (WHO) reports that 8 million people die each year from tobacco-related diseases. In 2019, 1.3 billion people were smoking, and 1.1 billion were using smokeless tobacco. Tobacco use is the leading preventable cause of death and disability worldwide.

#### 1. Lung cancer

Lung cancer is the leading cause of cancer death in both men and women. It is caused by the damage to the lungs from the tar and nicotine in tobacco smoke.

#### 2. Colorectal cancer

Colorectal cancer is the second leading cause of cancer death in both men and women. It is caused by the damage to the colon and rectum from the tar and nicotine in tobacco smoke.

#### 3. Stomach cancer

Stomach cancer is the fifth leading cause of cancer death in both men and women. It is caused by the damage to the stomach from the tar and nicotine in tobacco smoke.

#### 4. Bladder cancer

Bladder cancer is the sixth leading cause of cancer death in both men and women. It is caused by the damage to the bladder from the tar and nicotine in tobacco smoke.

#### 5. Pancreatic cancer

Pancreatic cancer is the eighth leading cause of cancer death in both men and women. It is caused by the damage to the pancreas from the tar and nicotine in tobacco smoke.

#### 6. Esophageal cancer

Esophageal cancer is the ninth leading cause of cancer death in both men and women. It is caused by the damage to the esophagus from the tar and nicotine in tobacco smoke.

#### 7. Liver cancer

Liver cancer is the tenth leading cause of cancer death in both men and women. It is caused by the damage to the liver from the tar and nicotine in tobacco smoke.

#### 8. Kidney cancer

Kidney cancer is the eleventh leading cause of cancer death in both men and women. It is caused by the damage to the kidneys from the tar and nicotine in tobacco smoke.

#### 9. Prostate cancer

Prostate cancer is the twelfth leading cause of cancer death in both men and women. It is caused by the damage to the prostate from the tar and nicotine in tobacco smoke.

#### 10. Cervical cancer

Cervical cancer is the thirteenth leading cause of cancer death in both men and women. It is caused by the damage to the cervix from the tar and nicotine in tobacco smoke.

#### 11. Breast cancer

Breast cancer is the fourteenth leading cause of cancer death in both men and women. It is caused by the damage to the breasts from the tar and nicotine in tobacco smoke.

#### 12. Ovarian cancer

Ovarian cancer is the fifteenth leading cause of cancer death in both men and women. It is caused by the damage to the ovaries from the tar and nicotine in tobacco smoke.

#### 13. Endometrial cancer

Endometrial cancer is the sixteenth leading cause of cancer death in both men and women. It is caused by the damage to the endometrium from the tar and nicotine in tobacco smoke.

#### 14. Thyroid cancer

Thyroid cancer is the seventeenth leading cause of cancer death in both men and women. It is caused by the damage to the thyroid from the tar and nicotine in tobacco smoke.

#### 15. Testicular cancer

Testicular cancer is the eighteenth leading cause of cancer death in both men and women. It is caused by the damage to the testicles from the tar and nicotine in tobacco smoke.

#### 16. Acute leukemia

Acute leukemia is the nineteenth leading cause of cancer death in both men and women. It is caused by the damage to the bone marrow from the tar and nicotine in tobacco smoke.

#### 17. Chronic leukemia

Chronic leukemia is the twentieth leading cause of cancer death in both men and women. It is caused by the damage to the bone marrow from the tar and nicotine in tobacco smoke.

#### 18. Multiple myeloma

Multiple myeloma is the twenty-first leading cause of cancer death in both men and women. It is caused by the damage to the bone marrow from the tar and nicotine in tobacco smoke.

#### 19. Glioblastoma

Glioblastoma is the twenty-second leading cause of cancer death in both men and women. It is caused by the damage to the brain from the tar and nicotine in tobacco smoke.

#### 20. Meningeal cancer

Meningeal cancer is the twenty-third leading cause of cancer death in both men and women. It is caused by the damage to the meninges from the tar and nicotine in tobacco smoke.

#### 21. Pancreatic cancer

Pancreatic cancer is the twenty-fourth leading cause of cancer death in both men and women. It is caused by the damage to the pancreas from the tar and nicotine in tobacco smoke.

#### 22. Stomach cancer

Stomach cancer is the twenty-fifth leading cause of cancer death in both men and women. It is caused by the damage to the stomach from the tar and nicotine in tobacco smoke.

#### 23. Liver cancer

Liver cancer is the twenty-sixth leading cause of cancer death in both men and women. It is caused by the damage to the liver from the tar and nicotine in tobacco smoke.

#### 24. Bladder cancer

Bladder cancer is the twenty-seventh leading cause of cancer death in both men and women. It is caused by the damage to the bladder from the tar and nicotine in tobacco smoke.

#### 25. Kidney cancer

Kidney cancer is the twenty-eighth leading cause of cancer death in both men and women. It is caused by the damage to the kidneys from the tar and nicotine in tobacco smoke.

#### 26. Prostate cancer

Prostate cancer is the twenty-ninth leading cause of cancer death in both men and women. It is caused by the damage to the prostate from the tar and nicotine in tobacco smoke.

#### 27. Breast cancer

Breast cancer is the thirtieth leading cause of cancer death in both men and women. It is caused by the damage to the breasts from the tar and nicotine in tobacco smoke.

#### 28. Ovarian cancer

Ovarian cancer is the thirty-first leading cause of cancer death in both men and women. It is caused by the damage to the ovaries from the tar and nicotine in tobacco smoke.

#### 29. Endometrial cancer

Endometrial cancer is the thirty-second leading cause of cancer death in both men and women. It is caused by the damage to the endometrium from the tar and nicotine in tobacco smoke.

#### 30. Thyroid cancer

Thyroid cancer is the thirty-third leading cause of cancer death in both men and women. It is caused by the damage to the thyroid from the tar and nicotine in tobacco smoke.

#### 31. Testicular cancer

Testicular cancer is the thirty-fourth leading cause of cancer death in both men and women. It is caused by the damage to the testicles from the tar and nicotine in tobacco smoke.

#### 32. Acute leukemia

Acute leukemia is the thirty-fifth leading cause of cancer death in both men and women. It is caused by the damage to the bone marrow from the tar and nicotine in tobacco smoke.

#### 33. Chronic leukemia

Chronic leukemia is the thirty-sixth leading cause of cancer death in both men and women. It is caused by the damage to the bone marrow from the tar and nicotine in tobacco smoke.

#### 34. Multiple myeloma

Multiple myeloma is the thirty-seventh leading cause of cancer death in both men and women. It is caused by the damage to the bone marrow from the tar and nicotine in tobacco smoke.

#### 35. Glioblastoma

Glioblastoma is the thirty-eighth leading cause of cancer death in both men and women. It is caused by the damage to the brain from the tar and nicotine in tobacco smoke.

#### 36. Meningeal cancer

Meningeal cancer is the thirty-ninth leading cause of cancer death in both men and women. It is caused by the damage to the meninges from the tar and nicotine in tobacco smoke.

#### 37. Pancreatic cancer

Pancreatic cancer is the fortieth leading cause of cancer death in both men and women. It is caused by the damage to the pancreas from the tar and nicotine in tobacco smoke.

#### 38. Stomach cancer

Stomach cancer is the forty-first leading cause of cancer death in both men and women. It is caused by the damage to the stomach from the tar and nicotine in tobacco smoke.

#### 39. Liver cancer

Liver cancer is the forty-second leading cause of cancer death in both men and women. It is caused by the damage to the liver from the tar and nicotine in tobacco smoke.

#### 40. Bladder cancer

Bladder cancer is the forty-third leading cause of cancer death in both men and women. It is caused by the damage to the bladder from the tar and nicotine in tobacco smoke.

#### 41. Kidney cancer

Kidney cancer is the forty-fourth leading cause of cancer death in both men and women. It is caused by the damage to the kidneys from the tar and nicotine in tobacco smoke.



[Cancer](#)

[Charge mondiale de morbidité](#)

[Éducation sanitaire](#)

[Épidémiologie](#)

[Femme](#)

[Législation sanitaire](#)

[Maladies cardiovasculaires](#)

Maladies non transmissibles, voir [Maladies chroniques](#)

[Promotion de la santé](#)

[Promotion de la santé à l'école](#)

[Surveillance de la santé publique](#)

[Tabac](#)

Saturday 17th of May 2025 04:27:49 PM