

Prevalence of allergenic arthropods in domestic dwellings of referrals to an asthma and allergy clinic in the Islamic Republic of Iran

Ayda Shafiei,¹ Mohammad D. Moemenbellah-Fard,² Kouros Azizi,² Seyed H. Nabavizadeh,³ Tahere Dabaghmanesh¹ and Aboozar Soltani²

¹Department of Medical Entomology and Vector Control; ²Research Center for Health Sciences, Institute of Health, Department of Medical Entomology and Vector Control, School of Health; ³Allergy Research Center, Department of Paediatrics, Shiraz University of Medical Sciences, Shiraz, Islamic Republic of Iran. (Correspondence to: Aboozar Soltani: abu2sol@yahoo.com).

Abstract

Background: Allergenic arthropods are crucial agents in inducing medically important respiratory diseases like asthma and the inflammation of the respiratory tract worldwide.

Aims: This study was conducted to determine the prevalence of all arthropods in the dwellings of people referred to the asthma and allergy clinic in Shiraz.

Methods: This was a cross-sectional descriptive study. Participants were 100 allergic patients who had tested positive (roach- and mite-sensitive). Mites were collected from their houses using a vacuum cleaner; other arthropods were caught with sticky traps. Direct observation and flotation methods were used and the samples were stored in 70% ethanol. Morphological characteristics were identified using valid taxonomic keys.

Results: Overall, 624 specimens were identified belonging to 14 orders (4 orders of mites: Astigmata, Cryptostigmata, Prostigmata and Mesostigmata; and 10 other arthropod orders: Diptera, Coleoptera, Hymenoptera, Thysanura, Thysanoptera, Entomobryomorpha, Blattodea, Siphonaptera, Pscoptera and Isopoda). The 2 most numerous species collected were *Musca domestica* and *Dermanyssus gallinae*.

Conclusion: A small number of dwellings were infested with roaches; none were infested with the common house dust mites. The allergies induced in these patients could likely be attributed to other arthropods that are not considered main allergens in asthma and allergy clinics in the Islamic Republic of Iran. Health surveillance and prevention of infestation for these arthropods could have an immense impact on the control of the allergenic arthropod community, prevention of respiratory diseases, and personal health in Shiraz.

Keywords: allergen, arthropod, asthma, allergy, indoor, urban

Citation: Shafiei A; Moemenbellah-Fard MD; Azizi K, Nabavizadeh SH; Dabaghmanesh T; Soltani A. Prevalence of allergenic arthropods in domestic dwellings of referrals to an asthma and allergy clinic in the Islamic Republic of Iran. *East Mediterr Health J.* 2019;25(x):xxx–xxx. <https://doi.org/10.26719/emhj.19.087>

Received: 23/05/18; accepted: 30/07/18

Copyright © World Health Organization (WHO) 2019. Some rights reserved. This work is available under the CC BY-NC-SA 3.0 IGO license <https://creativecommons.org/licenses/by-nc-sa/3.0/igo>

Introduction

Background

Over a million species of arthropods are reported in nature as having a significant role in the ecosystem; only a small fraction of these species are linked to human health. Arthropod vectors of infectious disease agents have special importance in the developing countries. They are also associated with certain allergic conditions and may induce or intensify some allergic reactions among humans (1,2).

In sensitive people, asthma symptoms may be triggered by inhaling allergens. The triggers differ from person to person, most likely due to a combination of environmental and genetic factors. Common asthma-causing allergens include: arthropods (dust mites, cockroaches, etc.), animal hair or dander, dust, respired chemicals, mould, pollen and tobacco smoke (1).

House dust mites are widespread, tiny (45–200 µm) arthropods belonging to the family Pyroglyphidae, order: Astigmata, class: Arachnida (3). These feed on skin relics, hair dandruff and other organic debris. They live and reproduce among clothes, bedding, carpets, furniture, household utilities and floors in human dwellings. The inhalation of faecal drops and protein moieties from the bodies of mites could cause allergic reactions like asthma, permanent allergic rhinitis and atopic dermatitis (eczema) among sensitive individuals (4,5).

Objectives

Many other arthropods are also considered or identified as being allergenic such as cockroaches, house flies, mosquitoes, storage pests, biting insects, canine and feline fleas, butterfly larvae, bedbugs, horseflies, silverfish and non-biting midges (Chironomidae). These could cause various types of allergy in some people (6). Any plan to control and prevent the spread of these arthropods must be based on their proper identification and classification. This study was thus conducted to gain some insight on species diversity of these allergenic arthropods, their abundance and extent of distribution in Shiraz.

Methods

Study area

This was a cross-sectional descriptive study conducted in Shiraz (29°40' N, 52°33' E), the capital city of Fars province, Islamic Republic of Iran. It is located at about 1500 m above sea level. It has a subtropical, hot, semi-arid climate with a mean annual temperature of 18 °C, relative humidity of 41% and precipitation 337.8 mm. Its hilly landscape is corrugated with the Zagros mountain range, which run northwest–southeast across the country.

Patients

In this investigation of all patients who were referred to the asthma and allergy clinic at Shiraz during the year of study, samplings were randomly carried out on the houses of 100 consenting patients who had positive skin tests to the *Dermatophagoides farinae* allergen, *D. pteronyssinus* allergen of house dust mites (*D. farinae*, *D. pteronyssinus*) and cockroach allergen. Samplings included patients' domiciles, in the particular loci which saw the highest frequency of inhabitants' activities. Participants were from both sexes with no age range limitation. After explaining the sampling procedure by an expert, informed consent forms were signed by all research participants or their parents (for children). In order to compensate for the possible drop-out of volunteers, sampling was continued until the sample size was completed.

Allergic skin reaction tests

Skin tests were done by the wheal and flare reactions on patients' forearms. Overall, 79 different items (comprising such categories as pollen, tree, grasses, weeds, mould, animals and food) were included in these tests, the reaction from each of which was observed directly on skin and the size of each reaction was measured and subsequently recorded. A drop of 50% glycerin + 50% COCAS fluid (containing NaCl 0.5%, NaHCO₃ 0.0275%, sterile water for injection, preservative 0.4% phenol) was used as negative control. A drop of standardized *D. pteronyssinus*, 5 mL in 10 000 arbitrary units/mL was used as the positive control. The negative control had a wheal and flare reaction of 1–3 units, while the positive control reaction measured > 3 units. They were compared 20 minutes after the initial administration of the drops and their sizes measured and recorded accordingly.

Sample size

The sample size was calculated according to the following formula:

$$n = \frac{(z_{1-\alpha} + z_{1-\beta})^2 p_0 q_0}{(p_1 - p_0)^2}$$

<>Sample size calculation for estimating a proportion p_1

Power	z-score	Precision	Variance	p_1	$p_1 \times (1-p_1)$
0.8	1.96	0.11	0.16	0.2	0.8
Sample size	100.7286				

Study design

At each house, samples were collected from different points using sticky traps or a vacuum cleaner to collect indoor arthropods. In the latter case, the content from each round of sampling (each 1m² in 1 minute) was emptied into a plastic freezer bag, labelled (collection place, relative humidity, ambient temperature, etc.) and frozen for later examination and identification in the laboratory. Sticky traps were fixed in different places and collected 24 hours later. The next day at the laboratory, equal weights of 200 mg portions from house dust samples were transferred into medium sized Petri dishes (minimum 5 times) and directly observed under a binocular microscope. Mites, being sensitive to light more slowly, were picked up and separated using the fine damp tip of entomological needle. They were then mounted onto microscope slides for further examination. Using diagnostic morphological features and valid taxonomic keys, they were identified to species level (3).

Statistical analysis

The collated data were uploaded onto the PC and analysed descriptively (frequencies and other descriptive statistics). We used *SPSS*, version 19, and *Excel* to tabulate results and draw histograms.

Results

A total of 624 specimens of arthropods were collected from the houses of 100 patients referred to the asthma and allergy clinic; these houses were located in various parts of Shiraz. From these, 46 mites and 578 other arthropods were identified. Table 1 shows the abundance order distribution of arthropods gathered from houses of patients referred to the asthma and allergy clinic. The species diversity in the insect orders and families was greater than in the class Arachnida. The specimens were classified into 14 orders based on their major morphological features. In the largest insect order, Coleoptera, 3 families, including 4 genera and 5 species, were identified (Table 1). In the medically important order Diptera, 4 families, including 4 species, were identified. Active search for arthropods using the vacuum cleaner method yielded a higher number of arthropod species (25 vs 16) than by the passive sticky trap collection method. The 2 most numerous species collected from patients' houses were *Musca domestica* and *Dermanyssus gallinae*. A total of 226 dipteran flies (129 *M. domestica*, 81 *Telmatoscopus proximus*, 11

Chrysomya albiceps, and 5 *Sarcodexia lambens*) were trapped from 7%, 20%, 3% and 2% of human residences, respectively (Table 1).

The moth fly, *T. proximus*, whose infestation by house was the highest among all diptera (20%) (Figure 1), has a pair of dichoptic eyes, clear venation on tapered wings, and 16-segmented digitiform antennae, each segment of which is proximally barrel-shaped and equipped with multiple rings of unbranched sensory filaments or ascoïd setae. This species is common in houses of Shiraz but is not considered an important allergenic arthropod by physicians or members of the health system. However, their hair and other part of their body could be allergenic to human.

The most numerous ($n = 97$) insect species found in patients' dwellings was the sand ant, *Myrmica sabuleti*, which was found in 24% of all infested houses (Figure 2). The second and third most frequent insect groups were the moth flies (mostly in suburban areas) and cockroaches (mostly in urban areas), being caught in 20% and 15% of all infested houses, respectively (Figure 1). All of the other insect species collected from patients' houses had infestation levels $\leq 10\%$.

The infestation frequencies of mites were more homogeneously restricted than insects, ranging from 1% to 5%, with the 2 species of red poultry mites (13 *Dermanyssus gallinae* and 6 *D. americanus*) being found in 5% of infested houses (Figure 3). The red poultry mite, *D. gallinae*, also had the highest rate (36.15%) of abundance among the mesostigmatid mites (Figure 4).

Discussion and conclusion

Our findings showed that all (100%) patients' dwellings were infested with at least one allergenic arthropod group. Generally, simultaneous multiple infestation of the dwellings of allergic human is the rule. Within the Arachnida, the cosmopolitan mesostigmatid red poultry mite, *D. gallinae*, represents a domestic and occupational allergenic species reported mostly from indoors in birds' nests. The allergenic character of this mite has been confirmed through clinical evidence as well as from purification and sequencing of amino acids in allergens. In 1970, Bernecker reported that 2% of individuals who were positive to the allergic dermal test from *D. pteronyssinus* were also positive to *D. gallinae*. The concept of immunological cross-reactivity from skin prick tests between these 2 different arthropod species should not be ignored. Allergic signs are thus indicated which could be induced by the bites of the red poultry mite. From this species, a tropomyosin allergen belonging to the group 10 allergens (*D. gallinae* allergen) has been isolated, cloned and sequenced (7). Cases of human infestation with this species were first reported in the Islamic Republic of Iran in 2014 (8). Some encephalitis viruses have been reported from this species, which is an indication of their medical importance. This species can also feed on humans, other mammals and birds and cause dermatitis and skin lesions. Cases of human infestation with *D. gallinae* are reported from Britain, Denmark, Egypt, France, Holland, Japan, Monte Negro, Morocco, Norway, Serbia and Turkey (9).

Isolation of the common rat house mite, *Haemolaelaps glasgowi*, from human dwellings could be attributed to the population eruption of urban rats infiltrating human dwellings in recent years. No routine antigenic tests are currently done for rat mites in this part of the world. Two other bird mites, *Ornithonyssus sylviarum* and *O. bursa*, which are often found among feathers, once on a human host could induce itching and painful dermatitis due to long allergic reactions. Some encephalitis viruses have been isolated from these 2 mites (10).

Description of *H. glasgowi* was first reported as the most widespread rodent ectoparasite in the Islamic Republic of Iran almost 2 decades ago (11). Both this species and *O. sylviarum* were subsequently identified from rodents in the city of Khoramabad in the west of the country (12). The latter species, the northern fowl mite, was recently described in sparrows in the west (13). The presence of these species was likely due to the breeding of either birds or rodents in human settlements in Shiraz.

The identification and medical importance of 3 other species of mesostigmatid mites, *Dermanyssus americanus*, *Laelaps nuttalli*, and *Echinolaelaps echidninus*, has been established in the Islamic Republic of Iran. All these species were reported as ectoparasites in a rodent control programme in the southern port city of Bandar-Abbas (14). The first of these, the American red poultry mite, often lives on birds. It can cause eczematous dermatitis in some individuals (15).

Only 2 species were found from the astigmatid mites, *Suidasia nesbitti* and *Glycyphagus prunorum*. Both of these have medical importance and could induce cross-reactivity with other allergenic mites. In addition, ingestion of foods infested with *S. nesbitti* could induce anaphylaxis in susceptible individuals (16). This species is known to be one of the group 2 allergen inducers. The cheese mite, *G. prunorum*, induces intense itching and pruritus by penetration into the fissures of the epidermal layers of foot, hand and face. It is most often found on workers in cereal, flour and vegetable storage facilities (17). This genus is considered to be a storage products and dust mite. Its allergens have been identified using the skin radioallergosorbent test (RAST), purification and amino acid sequencing. Cross-reactivity between this, storage, and house dust mites has previously been verified (18).

From the hexapods, beetles, including *Tribolium confusum*, were found, which are associated with rhinitis, conjunctivitis, wheal and signs of asthma in some individuals (19). The beetle genus *Dichillus* was first reported from Turkey in 2012 (20). No medical importance has so far been recorded for this genus.

From the order Hymenoptera, the sand ant, *Myrmica sabuleti*, was the most abundant species among the non-dipteran hexapods. In addition to fire ants, *Solenopsis*, other genera of ants like *Formica* and *Myrmica* could induce serious allergic reactions in people (21). The species *Vespula*

maculifrons, the eastern yellow jacket, is of medical importance but has not been reported from the Islamic Republic of Iran so far.

The order Diptera includes house flies, *Musca domestica* being the most frequently captured insect in allergic patients' dwellings. Their medical significance is evidently recorded. This and a number of other flies are also implicated in forensic medicine. The flesh fly, *Sarcodexia lambens*, may be involved in myiasis (22). Allergic reactions and asthma, particularly among children, may result from dried faeces, somatic setae, secretions and saliva from these species of flies in the vicinity of victims (19).

Steak bugs, including German and American roaches, *Blattella germanica* and *Periplaneta americana*, are 2 of the primary sources of allergens indoors inducing asthma and allergic reactions in susceptible individuals (23). The allergenic role and clinical importance of roaches on Iranian children with asthma has previously been investigated (24).

At present, studies on the allergic skin reaction test, which is routinely implemented in relation to allergenic arthropods at asthma and allergy clinics, is restricted to house dust mites (*D. farinae* and *D. pteronyssinus*) and cockroaches. Data from our investigation show that although all individuals whose houses were explored for the presence of arthropods were positive for the allergic skin reaction test, our findings indicate that only 15% of dwellings were infested with roaches and none of them were infested with the common house dust mites. Considering the fact that most arthropods can cross-react with other allergenic species, the allergies induced in these people could likely be attributed to other less important arthropods. It is thus suggested that, in line with routine tests, certain standard tests be carried out with regard to other allergenic arthropods so that the main disease-causing agent can be identified and suitable control measures can be included in an integrated control programme for training patients.

Finally, since a number of parasitic and vector-borne diseases are endemic to our region, the collection of data on the numerous prevalent arthropods is beneficial to the planning and evaluation of control measures to reduce disease. In conclusion, it is suggested that in all of the patients' dwellings, infestation with a minimal number of allergenic arthropod species are evidently observed. Health surveillance and prevention of infestation with these arthropods could have an immense impact on the control of the allergenic arthropod community, the prevention of respiratory diseases, and personal health care.

Acknowledgements

The authors are grateful for the assistance of the Vice-Chancellorship for Research and Technology at Shiraz University of Medical Sciences and to Mr H. Karami for his vigilant technical support.

Funding: This paper has partly been derived from, and supported by, the MSc thesis of Ms Ayda Shafiei (Grant N^o: 94-7574 dated 11 September 2015) in Medical Entomology, at the School of Health, Shiraz University of Medical Sciences, Islamic Republic of Iran.

Competing interests: None declared.

References

1. Jeong KY, Hong CS, Yong TS. Domestic arthropods and their allergens. *Protein Pept Lett*. 2007 Oct 1;14(10):934–42. PMID:18220990
2. Linneberg A. Are we getting enough allergens? *Int Arch Allergy Appl Immunol*. 2008 Jun 3;147(2):93–100. doi:10.1159/000135695
3. Service MW. *Medical entomology for students*, 3rd ed. Cambridge: Cambridge University Press; 2004.
4. Wang DY. Risk factors of allergic rhinitis: genetic or environmental. *Ther Clin Risk Manag*. 2005 Jun;1(2):115–23. PMID:18360551
5. Soltani A, Azizi K, Saleh V, Dabaghmanesh T. The fauna and distribution of house dust mites in residential homes of Bandar Abbas District, Southern Iran. *Exp Appl Acarol*. 2011 Jul 1;54(3):269–76. doi:10.1007/s10493-011-9436-6
6. Arlian LG. Arthropod allergens and human health. *Annu Rev Entomol*. 2002 Jan;47(1):395–433. PMID:11729080
7. Nisbet AJ, Huntley JF, Mackellar A, Sparks N, McDevitt R. A house dust mite allergen homologue from poultry red mite *Dermanyssus gallinae* (De Geer). *Parasite Immunol*. 2006 Aug 1;28(8):401–5. doi:10.1111/j.1365-3024.2006.00862.x
8. Abdigoudarzi M, Mirafzali MS, Belgheiszadeh H. Human infestation with *Dermanyssus gallinae* (Acari: Dermanyssidae) in a family referred with pruritus and skin lesions. *Iran J Arthropod Borne Dis*. 2014 Jun 1;8(1):119. PMID:25629073
9. Dogramaci AC, Culha G, Özçelik S. *Dermanyssus gallinae* infestation: an unusual cause of scalp pruritus treated with permethrin shampoo. [J Dermatolog Treat](#). 2010 Sep;21(5):319–21. doi:10.3109/09546630903287437
10. Colloff MJ. *Dust mites*. Dordrecht: Springer; 2009. doi:10.1007/978-90-481-2224-0
11. Bochkov A, Arbobi M, Malikov V. Notes on mites of the family Myobiidae (Acari: Prostigmata) parasitising rodents (Mammalia: Rodentia) in Iran. *Folia Parasitol*. 2000 Jan 1;47(1):73–7. PMID:10833020

12. Shayan A, Rafinejad J. Arthropod parasites of rodents in Khorram Abbad district, Lorestan Provincen of Iran. *Iran J Public Health*. 2006;35(3):70–6.
13. Moodi B, Aliabadian M, Moshaverinia A, Kakhki OM, Faraji F. Mites associated with passerine birds in eastern Iran. *Int J Acarology*. 2014 Feb 17;40(2):133–7. doi:10.1080/01647954.2014.888094
14. Hanafi-Bojd AA, Shahi M, Baghaili M, Shayeghi M, Razmand N, Pakari A. A study on rodent ectoparasites in Bandar Abbas: the main economic southern seaport of Iran. *Iran J Environ Health Sci Eng*. 2007;4(3):173–6.
15. George DR, Finn RD, Graham KM, Mul MF, Maurer V, Moro CV, et al. Should the poultry red mite *Dermanyssus gallinae* be of wider concern for veterinary and medical science?. *Parasit Vectors*. 2015 Mar 25;8(1):1. doi:10.1186/s13071-015-0768-7
16. Dutau G. House dust mites: new food allergens. *Rev Francaise d'Allergologie et d'Immunologie Clinique*. 2002 Mar 1;42(2):171–7.
17. Steen CJ, Carbonaro PA, Schwartz RA. Arthropods in dermatology. *J Am Acad Dermatol*. 2004 Jun 30;50(6):819–42. doi:10.1016/j.jaad.2003.12.019
18. Mumcuoglu Y. House dust mites in Switzerland: II. culture and control. *Int J Acarol*. 1977 Jun 1;3(1):19–25. PMID:1162310
19. Arlian LG. Arthropod allergens and human health. *Annu Rev Entomol*. 2002 Jan;47(1):395–433. PMID:11729080
20. Canpolat D, Hasbenli A. New records of Tenebrioninae and Pimeliinae (Coleoptera: Tenebrionidae) from Turkey. *J Entomol Res Soc*. 2012 Jan 1;14(1):15–20.
21. Horton SM. Identifying the locations, movement and habitat of the European fire ant, *Myrmica rubra*: an invasive species in the urban/suburban environment of Halifax, Nova Scotia [thesis]. Halifax: Saint Mary's University; 2012.
22. Bermúdez C, Buenaventura R, Couri M, Miranda RJ, Herrera JM. Mixed myiasis by *Philornis glaucinis* (Diptera: Muscidae), *Sarcodexia lambens* (Diptera: Sarcophagidae) and *Lucilia eximia* (Diptera: Calliphoridae) in *Ramphocelus dimidiatus* (Aves: Thraupidae) chicks in Panama. *Bol Soc Entomol Aragon*. 2010(47):445–6.
23. Mollet JA, Vailes LD, Avner DB, Perzanowski MS, Arruda LK, Chapman MD, et al. Evaluation of German cockroach (Orthoptera: Blattellidae) allergen and seasonal variation in low-income housing. *J Med Entomol*. 1997 May 1;34(3):307–11. doi:10.1093/jmedent/34.3.307
24. Farhoudi A, Pourpak Z, Mesdaghi M, Chavoshzadeh Z, Kazemnejad A. The study of cockroach allergy in Iranian children with asthma. *Iran J Med Sci*. 2015 Nov 30;27(4):156–60.

Table 1. Abundance (%) order distribution of arthropods collected from houses of patients referred to the asthma and allergy clinic of Shiraz

Order	Family	Genus	Species	Common name	No. trapped	Abundance (%)	How collected
Insecta							
Collembola	Entomobryidae	Entomobrya	multifasciata	Spring tail	45	90	VC
			nicoletti	Spring tail	5	10	VC
Coleoptera	Tenebrionidae	Tribolium	castaneum	Red flour beetle	28	36.36	VC/ST
			confusum	Confused flour beetle	10	13	VC/ST
		Dichillus	Unidentified	Darkling beetle	8	10.38	VC
	Cucujidae	Oryzaephilus	surinamensis	Saw-toothed grain beetle	4	5.19	VC/ST
	Dermeestidae	Anthrenus	museorum	Museum/skin beetle	5/22	35.07	VC/ST
Hymenoptera	Formicidae	Myrmica	sabuleti	Sand ant	97	92.38	VC/ST
	Vespidae	Vespula	macalifrons	Eastern yellow jacket	2	1.91	ST
		Polistes	gallicus	Paper wasp	6	5.71	ST
Diptera	Muscidae	Musca	domestica	House fly	129	57.07	ST
	Calliphoridae	Chrysomya	albiceps	Blow fly	11	4.87	ST
	Sarcophagidae	Sarcodexia	lambens	Flesh fly	5	2.22	ST
	Psychodidae	Telmatoscopus	proximus	Moth fly	81	35.84	ST
Blattodea	Blattidae	Periplaneta	americana	American cockroach	12	43.55	ST
	Blatellidae	Blatella	germanica	German cockroach	20	56.45	ST
Thysanura	Lepismatidae	Lepisma	saccharina	Silver fish	15	100	VC
Thysanoptera	Thripidae	Frankliniella	tritici	Flower thrips	16	100	VC
Siphonaptera	Pulicidae	Pulex	irritans	Human flea	1	100	ST
Psocoptera	Liposcelididae	Liposcelis	paetus	Book louse	10	100	VC/ST
Malacostraca							

Isopoda	Cylisticidae	Cylisticus	convexus	Wood louse	16	100	VC/ST
Arachnida: Acarid mites							
Astigmata	Glycyphagidae	Glycyphagus	prunorum	Cheese mite	1	25	VC
	Suidasiidae	Suidasia	nesbitti	Scaly grain mite	3	75	VC
Cryptostigmat	Oppiidae	Aeroppia	SU	Oribatid mite	5	100	VC
a							
Prostigmata	Bdellidae	Cyta	latirostris	Predator mite	1	100	VC
Mesostigmata	Laelaptidae	Haemolaelaps	glasgowi	Common rodent mite	3	8.33	VC
		Echinolaelaps	echidninus	Spiny rat mite	4	11.11	VC
		Laelaps	nutalli	Domestic rat mite	3	8.33	VC
	Dermanyssidae	Dermanyssus	gallinae	Red fowl mite	13	36.15	VC
			americanus	American bird mite	6	16.66	VC
	Macronyssidae	Ornithonyssus	bursa	Tropical bird mite	3	8.33	VC
			sylviarum	Northern fowl mite	2	5.55	VC
	Ascidae	Blattisocius	tarsalis	Egg eating mite	1	2.77	VC
	Parasitidae	Holoparasitus	Unidentified	Gamasid mite	1	2.77	VC

VC = vacuum cleaner; ST = sticky trap.




Figure 1. Distribution of infestations of diptera, fleas and cockroaches in allergic patients' dwellings ($n = 100$) in Shiraz, 2016




Figure 2. Distribution of infestations of non-dipteran insects found in allergic patients' dwellings (n = 100) in Shiraz, 2016




Figure 3. Distribution of all mites found in allergic patients' dwellings ($n = 100$) in Shiraz, 2016




Figure 4. Distribution of infestations of mesostigmatid mites found in allergic patients' dwellings, Shiraz, 2016