

## **Workplace violence among health care workers in emergency departments of public hospitals in Dammam, Saudi Arabia**

Moussa M. Harthi<sup>1</sup>, Mohammed A. Olayan<sup>1</sup>, Hassan A. Abugad<sup>1</sup> and Moataza M. Abdel Wahab<sup>1</sup>

<sup>1</sup>Department of Family and Community Medicine, Imam Abdulrahman Bin Faisal University, Dammam, Saudi Arabia (Correspondence to: M.M. Harthi: dr\_mmh@hotmail.com).

### **Abstract**

**Background:** Workplace violence is a serious occupational health problem. Emergency healthcare workers (HCWs) have a high risk of exposure to violence with negative personal consequences.

**Aims:** To estimate the prevalence and possible associated factors of workplace violence among HCWs in emergency departments (EDs) of public hospitals in Dammam, Saudi Arabia.

**Methods:** A cross-sectional study was conducted during August to October 2018 at 4 EDs of public hospitals belonging to the Saudi Ministry of Health. Data were collected using a self-administered questionnaire.

**Results:** Of 380 questionnaires distributed, 324 were returned (85% response rate). Almost two thirds of the participants were women (66.4%) and more than half (54%) were nurses. A total of 155 HCWs (47.8%) had experienced at least 1 type of violent incident in the preceding 12 months. Of the total violence incidents, 52% were verbal abuse, 19% were physical violence, and sexual harassment (3%) was the least common. Lack of encouragement to report incidents and Saudi nationality were the only significant variables associated with workplace violence.

**Conclusions:** Workplace violence was prevalent, and verbal abuse was the commonest type among HCWs in emergency departments of Saudi hospitals. Encouragement to report violent incidents and raising awareness among HCWs about violence reporting systems are important strategies to improve workplace safety.

**Keywords:** emergency departments, healthcare workers, Saudi Arabia, reporting, workplace violence

Citation: Harthi MM; Olayan MA; Abugad HA; Abdel Wahab MM. Workplace violence among health care workers in emergency departments of public hospitals in Dammam, Saudi Arabia. East Mediterr Health J. 2020;26(x):xxx-xxx <http://doi.org/10.26719/emhj.20.069>

Received: 30/05/19; accepted: 27/11/19

## Introduction

Health care workers (HCWs) are among the groups most experiencing violence and aggressive behaviour at work, especially those who work in emergency departments (EDs) in public hospitals (1). Workplace violence has negative consequences on safety and workplace activities of HCWs (2). However, the estimated prevalence of violence against HCWs is still unknown because there is no clear definition of a violent incident (1,2). The World Health Organization (WHO) defined violence as “The intentional use of physical force or power, threatened or actual, against another person or against oneself or a group of people that results in or has a high likelihood of resulting in injury, death, psychological harm, maldevelopment or deprivation” (3). The National Institute for Occupational Safety and Health defines workplace violence as “violent acts (including physical assault and threats of assault) directed towards persons at work or on duty” (4). According to WHO, physical or psychological violence can appear in different forms, which may often overlap (4,5). Physical violence is defined as the use of physical force against another person or group that results in physical, sexual or psychological harm, and such violence includes beating, kicking, slapping, stabbing, shooting, pushing, biting and pinching (3,5). Psychological violence is defined as intentional use of power, including threat of physical force, against another person or group that can result in harm to physical, mental, spiritual, moral or social development. Psychological violence includes verbal abuse, bullying/mobbing, harassment (including sexual and racial) and threats.

Many studies worldwide have examined the prevalence of workplace violence among HCWs (2). A survey of workplace violence across 65 American EDs conducted in 2008 showed that the violence and weapons in the EDs were common, and nurses were less likely to feel safe than other staff were (6). A cross-sectional study in 2009 in Tokyo, Japan revealed that 36.4% of 11 095 HCWs in 19 hospitals experienced workplace violence by patients or their relatives; 15.9% experienced physical aggression, 29.8% experienced verbal abuse and 9.9% experienced sexual harassment (7). In another large study conducted between October 2012 and July 2013 at primary healthcare centres in Belgrade, Serbia, the prevalence of workplace violence was 52.6% among 1757 HCWs (8). In the Middle East, workplace violence has been investigated in several studies. An Iranian cross-sectional survey in 2011 among 196 nurses in 11 EDs in teaching hospitals in Tehran, showed that 19.7% of nurses faced physical violence and 91.6% experienced verbal abuse (9). Another cross-sectional study in Jordan in 2011 among 227 nurses in 12 provinces revealed that 75.8% were exposed to at least 1 type of violence (10). A comprehensive survey of workplace violence among 713 physicians in EDs in Turkey found that 78.1% had experienced violence (11).

Factors related to the increased the risk of workplace violence are related to the offenders, HCWs or the workplace environment (2). Personality and mental health disorders (such as schizophrenia, paranoia, anxiety, antisocial attitude, dementia and alcohol abuse) are the most significant factors related to the offenders (7). HCW-related factors include understaffed working conditions, working alone and long working hours (7,12). Factors related to the workplace include long waiting times, overcrowding, inadequate security, and lack of policies for preventing violence (12).

In a few studies in Saudi Arabia, there was difficulty in estimating the magnitude of the problem due to lack of reporting and other factors (2,13). In 2009, a self-reporting questionnaire study in Al-Hassa of 1091 primary health care professionals revealed that 28% suffered from workplace violence (12). A cross-sectional study in Riyadh in 2011 of 600 physicians and nurses found that 67.4% were exposed to workplace violence, and that nurses were more susceptible than physicians (14). In another cross-sectional study in 2014 in 12 family medical centres in Riyadh, 45.6% of 270 HCWs experienced some sort of violence during the 12 months prior to the study (2). Three studies were conducted in Saudi Arabia in 2015. A cross-sectional study at King Fahd Hospital showed that 30.7% of 391 nurses were exposed to verbal abuse (13). In EDs of 3 hospitals in Riyadh, 89.3% of 121 nurses experienced a violent incident in the 12 months prior to the study (15). In EDs in Tabuk, 90.7% of 129 had history of workplace violence (1). EDs are in operation 24 hours a day, 7 days a week (16). Patients usually come to EDs with relatives or friends with expectations of a rapid response and good service from HCWs regardless of the severity of the case (12). EDs receive a huge number of patients, therefore, the chance of HCWs being exposed to violence is high (1,12).

This study was conducted to estimate the prevalence of workplace violence among HCWs in EDs in public hospitals in Dammam, Saudi Arabia and to determine possible associated factors.

## **Methods**

This was a cross-sectional survey conducted during August to October 2018 at 4 public hospitals belonging to the Ministry of Health in Dammam, Saudi Arabia: Dammam General Medical Complex, Dhahran Eye Specialist Hospital, Maternal and Children's Hospital and Al-Amal Complex for Mental Health). All HCWs in all duty shifts (morning, evening and night) in EDs were invited to participate, with exclusion of those with work experience < 1 year. The sample size was calculated using epi info, assuming the level of violence among HCWs was 89% from previous data (15), with an accepted margin of error 4%. The sample by population survey was 235 HCWs at 95% confidence level and was increased to 294 HCWs, expecting 80% response.

Data were collected from 324 participants, using a self-administered questionnaire that was based on questionnaires developed by WHO (5) and was modified by the researchers. The

English language questionnaire was translated into Arabic by the authors and validated by 3 experts in the Department of Family and Community Medicine, Imam Abdulrahman Bin Faisal University. The questionnaire consisted of 8 sections. The first part included demographic information such as age, sex, marital status, occupational title, nationality, educational level, and years of work experience. The second part consisted of items that addressed occupational characteristics (working multiple shifts, shift time worked, number of coworkers in the same work area, encouragement to report violent events, and availability of a violence reporting system). The other sections consisted of items that addressed the characteristics of the violent acts experienced (time, place and frequency of violence) and the identity, age and sex of the offender. There were also questions about reasons for violence (e.g., lack of security and absence of punishment) and the consequences for the HCWs and the offenders. Finally, there was a question about reasons for not reporting acts of violence. Types of violence were classified into physical, verbal, bullying, and sexual and racial harassment (5). A pilot study was carried out on 10 HCWs in 1 public hospital on 1 day, to check the clarity of the language used and estimate the average time to answer the questionnaire. The participants in the pilot study were not included in the present study.

All statistical analyses were conducted using SPSS version 25, setting our level of confidence at 95%. Descriptive statistics by frequency and percentage were used for categorical variables, while continuous variables were assessed for normality. The frequency of workplace violence was calculated by dividing the number of those who had experienced violence during the preceding 12 months by the total number of HCWs in the study. The  $\chi^2$  and independent samples *t* test were used to assess the relation between demographic and occupational characteristics and workplace violence. Logistic regression analysis was used to assess factors independently associated with the occurrence of workplace violence. Adjusted odds ratios with corresponding 95% confidence intervals were presented.

## **Results**

### *Demographic and occupational characteristics*

Of 380 questionnaires distributed, 324 were returned (85% response rate). The age of participants ranged between 22 and 55 years, with a mean of 32.7 (standard deviation, 6.2) years, and 215 were women (66.4%) (Table 1). The majority (78.1%) of HCWs were Saudis and almost two thirds were married. The largest proportion had a diploma (50.3%) followed by a bachelor's degree (43.5%). More than half the HCWs (54%) were nurses and 40.1% had work experience of 6–9 years.

### *Report encouragement and system availability*

One hundred and ninety-three (59.6%) of 324 respondents stated that they were encouraged to report workplace violence and 131 (40.4%) that they were not encouraged. Two hundred and twenty-five (69.4%) HCWs reported that a system was available for reporting violence and 99 (30.6%) reported no such system.

#### *Frequency and type of violent incident*

Out of 324 HCWs, 155 (47.8 %) had experienced at least 1 type of violent incident during the preceding 12 months. Among 241 incidents, 126 (52%) were verbal abuse, 45 (19%) physical violence, 39 (16%) bullying, 24 (10%) racial harassment and 7 (3%) sexual harassment (Table 2). Ninety-five (39.4%) violent incidents happened in the morning and the same number in the evening. Almost all (n = 232, 96.3%) of the violent incidents occurred in the workplace. Ninety-nine (41.1%) violent incidents occurred once a year and 73 (30.3%) more than once a month. Most (n = 102, 42%) of the offenders were patients, followed by relatives of patients (n = 75, 31%). The majority (n = 197, 82%) of the offenders were aged 21–45 years and 41 (17%) were ≥ 46 years. Both men and women committed the violent act in 97 (40.25%) cases, men only in 95 (39.42%) and women only in 49 (20.33%). Most (n = 180, 74.7%) of the participants exposed to violence believed that the incident could have been prevented. The violence incident ended with the following consequences for the offenders: none (n = 154, 63.9%), verbal warning (n = 51, 21.2%) and reported to the police (n = 16, 6.6%). The consequences for HCWs were: none (n = 112, 46.5%), reduced work performance (n = 107, 44.4%), documented complaint against HCWs (n = 20, 8.3%) and injuries (n = 2, 0.8%). Almost all (n = 41, 91.1%) incidents of physical violence happened without a weapon and 23 (51.1%) were committed by men. Most physical (n = 22, 48.9%) and verbal abuse (n = 55, 43.7%) occurred in the evening. The majority (n = 29, 74.4%) of bullying incidents occurred in the morning and managers were a major source (n = 22, 56.4%) of violent incidents, followed by staff members (n = 14, 35.9%). In 34 (87.2%) of those incidents, no action was taken. Sexual harassment among staff members was the highest (n = 3, 42.9%). Decline in work performance was reported in 20 (44.4%) HCWs who experienced physical violence and in 13 (54.2%) who were subjected to racial harassment.

#### *Factors associated with workplace violence*

HCWs who experienced violence reported that it was caused by absence of punishment (67%), lack of security (51%), staff shortage (34%), long waiting time for patients (33%), overcrowding (29%), personality type (17%), cultural beliefs (9%), lack of patient privacy (3%) and language barrier (2%). Absence of punishment was the most common cause of verbal abuse (61%), bullying (95%), sexual (71%) and racial (58%) harassment, whereas, lack of security was the most common cause of physical violence (64%).

#### *History of workplace violence related to characteristics of HCWs*

Demographic and occupational features of HCWs who did and did not experience violence are shown in Table 3. Sex was significantly associated with violence, with violence being more frequent for men (n = 63, 57.8%) than women (n = 92, 42.8%). Nationality was significantly associated with violence and was more frequent for Saudis (n = 131, 51.8%) than non-Saudis (n = 24, 33.8%). Those who worked with ≤ 10 coworkers (n = 124, 53.4%) reported significantly more frequent violence than those who worked with > 10 coworkers (n = 31, 33.7%). Those

who lacked encouragement to report violent acts (n = 79, 60.3%) reported significantly more frequent violence than those who had such encouragement (n = 76, 39.4%). Those who confirmed lack of availability of a system for reporting violence (n = 57, 57.6%) reported significantly more frequent violence than those who confirmed system availability (n = 98, 43.6%).

#### *Type of workplace violence related to characteristics of HCWs*

Men (n = 22, 20.2%) experienced significantly more physical violence than women did (n = 23, 10.7%) (Table 4). Men (n = 51, 46.8%) also had significantly more verbal abuse than women had (n = 75, 34.9%). Violence was significantly more frequent for unmarried (n = 5, 5.1%) than married (n = 2, 0.9%) HCWs. Saudi HCWs (n = 106, 41.9%) experienced verbal abuse significantly more often than non-Saudis did (n = 20, 28.2%). Physical violence was significantly more frequent in HCWs with < 10 coworkers (n = 38, 16.4%) than in those with > 10 coworkers (n = 7, 7.6%). Verbal abuse was also significantly more frequent in HCWs with < 10 coworkers (n = 101, 43.5%) than in those with > 10 coworkers (n = 25, 27.2%). HCWs who lacked encouragement to report violent incidents reported significantly more verbal abuse (n = 65, 49.6%) than those who had encouragement (n = 61, 31.6%). Similarly, HCWs who lacked encouragement to report violence reported significantly more bullying (n = 23, 17.6%) than those who had encouragement (n = 16, 8.3%). In contrast, demographic and occupational characteristics, such as age, occupation, shift time, direct contact with patient, and patient types, were not significantly associated with general or specific types of violence.

#### *Logistic regression analysis of workplace violence*

After entering sex, nationality, number of coworkers, lack of report encouragement and system availability into the regression model, the only independent variables significantly associated with general violence were lack of report encouragement and Saudi nationality (Table 5). For physical violence, the only significant independent factor was male sex. Lack of report encouragement was the only variable that remained significantly associated with verbal abuse and bullying.

### **Discussion**

The main aim of this study was to estimate the prevalence of workplace violence in a sample of 324 participants working in EDs in 4 public hospitals in Dammam, Saudi Arabia. The study showed that the prevalence of violence among HCWs was 47.8%, which was considerably lower than 89.3% in nurses in the EDs in 3 public hospitals in Saudi Arabia (15). However, our result was closer to the prevalence of 57.5% in HCWs in 2 government hospitals and 10 primary healthcare centres in Saudi Arabia who experienced at least 1 violence incident (22), and similar to the prevalence of 45.6% among HCWs in 12 family medical centres in Riyadh (2).

Most studies have shown that psychological violence (especially verbal abuse) was higher than physical violence (15,21,26). The number of incidents of verbal abuse was approximately 5-fold that of the number of incidents of physical violence among nurses in several EDs in Jordan (10), which can be explained by the stress of acute illness experienced by patients and/or families at the time of the violent act. In the current study, verbal abuse formed 52% of the violent incidents, physical violence 19%, bullying 16%, racial harassment 10% and sexual harassment was the least common (3%). Similarly, a study in Macau revealed incidents of verbal abuse (53.4%), physical assault (16.1%), bullying (14.2%), sexual harassment (4.6%) and racial harassment (2.6%) among physicians and nurses (24). Verbal abuse was the most common form of violence because it was easy to perpetuate and could not be controlled by any sort of security measures. The majority (74.4%) of bullying incidents occurred in the morning and managers were a major source (56.7%) of incidents followed by staff members (35.9%), which is often explained by the presence of most managerial staff in the morning. Moreover, interprofessional violence may have played a role in these incidents.

Most of the workplace violence was experienced by Saudi nationals, which is explained mainly by the high number of Saudi participants in the study. The majority of offenders were patients (42%) followed by their relatives (31%), which was similar to some previous studies (2,15, 20,25,26) but contrary to others (1,14,23), in which the companions of the patients were the main offenders. The fact that patients were the major aggressors in the current study could be explained by the absence of deterrent action (63.9%) towards violent incidents as supported by management in the workplaces, following the rule “the patient is always right”.

Workplace violence had negative consequences on HCWs, such as reduced work performance (44.4%), complaints against HCWs (8.3%) and injuries (0.8%), which is supported by previous studies (1,2,18,19). Reduced work performance could be explained by feeling unsafe, anger, anxiety or distress or performing duties in an unprofessional way. Some previous studies suggested that the reasons for violence in EDs were staff shortage, absence of punishment, lack of security, and long waiting times for patients. Certain characteristics of HCWs, including age, sex, years of experience and marital status, have been associated with increased workplace violence (27,28). In the current study, the frequency of physical violence was high among men (20.2%) and sexual harassment was high among unmarried HCWs (5.1%).

More than half of violent incidents (66.7 %) were not reported and the main reason was the feeling that reporting was useless. This could be related to the existing system that includes reporting the incident to a supervisor, duty director, or the police. Most HCWs (n = 57, 57.6%) exposed to workplace violence questioned the availability of a violence reporting system. Moreover, the majority (75.9%) raised queries about the efficiency of the security measures applied in EDs of the studied hospitals. Our results could be explained by lack of awareness of the reporting systems and inefficient security measures. Hogarth et al. (29) noted that the

solution agreed upon by HCWs to decrease workplace violence was encouragement by management to report violent incidents and to develop preventative measures.

The current study is one of few to cover all types of violence (physical and psychological, including verbal threats, bullying, and sexual and racial harassment) and used the standard WHO definition of violence. Additionally, all HCWs in EDs of public hospitals were targeted. However, limitations cannot be excluded. The size of the sample may limit generalization of the results. The questionnaire was self-administered and recall bias could not be excluded, as in most similar surveys.

## Conclusions

In this study, workplace violence was prevalent among HCWs, and verbal abuse was the commonest type. The most important associated factor was absence of punishment, which was agreed upon by the majority of HCWs. Creation of an environment that encourages HCWs to report violent incidents and raising awareness of HCWs about violence reporting systems in EDs are recommended. Ensuring the reporting of all violent incidents and follow-up of the appropriate actions are essential. Supporting programmes to help and provide HCWs with the knowledge to manage and control incidents are needed.

## References

- (1) Alzahrani TY, Almutairi AH, Alamri DA, Alamri MM and Alalawi YS. Violence and aggression toward health care professionals in emergency departments in Tabuk, Saudi Arabia. *Eur J Pharm Med Res*. 2016,3(1):5–11.  
<http://www.ejpmr.com/admin/download/article/MTQ1MTUzODA3NS5wZGY>
- (2) Al-Turki N, Afify AA and AlAteeq M. Violence against health workers in Family Medicine Centers. *JMDH*. 2016 May 31;9:257–66.  
<https://www.ncbi.nlm.nih.gov/pubmed/27330300> PMID:27330300
- (3) Krug EG, Mercy JA, Dahlberg LL and Zwi AB. The world report on violence and health. *The Lancet*. 2002 Oct 5;360(9339):1083–8  
<https://www.ncbi.nlm.nih.gov/pubmed/12384003> PMID:12384003
- (4) Violence. Occupational hazards in hospitals. Cincinnati: National Institute for Occupational Safety and Health; 2002 (<https://www.cdc.gov/niosh/docs/2002-101/default.html>, accessed 6 May 2020).
- (5) ILO, WHO. Joint program on workplace violence in the health sector, workplace violence in the health sector. Country case studies research instruments. Survey questionnaire. English. Geneva: International Labour Office, International Council of Nurses, World Health Organization, Public Services International; 2003  
([https://www.who.int/violence\\_injury\\_prevention/violence/interpersonal/en/WVquestionnaire.pdf](https://www.who.int/violence_injury_prevention/violence/interpersonal/en/WVquestionnaire.pdf), accessed 6 May 2020).

- (6) Kansagra SM, Rao SR, Sullivan AF, Gordon JA, Magid DJ, Kaushal R, et al. A survey of workplace violence across 65 US emergency departments. *Acad Emerg Med*. 2008 Dec;15(12):1268–74. <http://dx.doi.org/10.1111/j.1553-2712.2008.00282.x> PMID:18976337
- (7) Fujita S, Ito S, Seto K, Kitazawa T, Matsumoto K and Hasegawa T. Risk factors of workplace violence at hospitals in Japan. *J Hosp Med*. 2012 Feb;7(2):79–84. <http://dx.doi.org/10.1002/jhm.976> PMID:22058040
- (8) Fisekovic MB, Trajkovic GZ, Bjegovic-Mikanovic VM and Terzic-Supic ZJ. Does workplace violence exist in primary health care? Evidence from Serbia. *Eur J Public Health*. 2015 Aug;25(4):693–8. <http://dx.doi.org/10.1093/eurpub/cku247> PMID:25644138
- (9) Esmaeilpour M, Salsali M and Ahmadi F. Workplace violence against Iranian nurses working in emergency departments. *Int Nurs Rev*. 2011 Mar;58(1):130–7. <http://dx.doi.org/10.1111/j.1466-7657.2010.00834.x> PMID:21281305
- (10) Albashtawy M. Workplace violence against nurses in emergency departments in Jordan. *Int Nurs Rev*. 2013 Dec 1;60(4):550–5. <http://dx.doi.org/10.1111/inr.12059> PMID:24117233
- (11) Bayram B, Çetin M, Oray NÇ and Can İÖ. Workplace violence against physicians in Turkey's emergency departments: a cross-sectional survey. *BMJ open*. 2017 Jun 29;7(6):e013568. <http://dx.doi.org/10.1136/bmjopen-2016-013568> PMID:28663323
- (12) El-Gilany AH, El-Wehady A and Amr M. Violence against primary health care workers in Al-Hassa, Saudi Arabia. *JIV*. 2010 Apr;25(4):716–34. <http://dx.doi.org/10.1177/0886260509334395> PMID:19494243
- (13) Al-Shamlan NA, Jayaseeli N, Al-Shawi MM and Al-Joudi AS. Are nurses verbally abused? A cross-sectional study of nurses at a university hospital, Eastern Province, Saudi Arabia. *J Family Community Med*. 2017 Sep–Dec;24(3):173–80. [http://dx.doi.org/10.4103/jfcm.JFCM\\_45\\_17](http://dx.doi.org/10.4103/jfcm.JFCM_45_17) PMID:28932162
- (14) Algwaiz WM and Alghanim SA. Violence exposure among health care professionals in Saudi public hospitals. A preliminary investigation. *Saudi Med J*. 2012 Jan;33(1):76–82. PMID:22273653
- (15) Alyaemni A and Alhudaithi H. Workplace violence against nurses in the emergency departments of three hospitals in Riyadh, Saudi Arabia: a cross-sectional survey. *NursingPlus Open*. 2016;2:35-41. <https://doi.org/10.1016/j.npls.2016.09.001>
- (16) Rehmani R. Emergency section and overcrowding in a university hospital of Karachi, Pakistan. *JPMMA*. 2004 May;54(5):233–7. PMID:15270179
- (17) Abdellah RF and Salama KM. Prevalence and risk factors of workplace violence against health care workers in emergency department in Ismailia, Egypt. *Pan Afr Med J*. 2017 Jan 17;26:21. <http://dx.doi.org/10.11604/pamj.2017.26.21.10837> PMID:28451000
- (18) Magnavita N and Heponiemi T. Violence towards health care workers in a Public Health Care Facility in Italy: a repeated cross-sectional study. *BMC Health Serv Res*. 2012Dec;12(1):108. <http://dx.doi.org/10.1186/1472-6963-12-108> PMID:22551645

- (19) Kitaneh M and Hamdan M. Workplace violence against physicians and nurses in Palestinian public hospitals: a cross-sectional study. *BMC Health Serv Res*. 2012 Dec;12(1):469.  
<https://bmchealthservres.biomedcentral.com/track/pdf/10.1186/1472-6963-12-469>
- (20) Speroni KG, Fitch T, Dawson E, Dugan L and Atherton M. Incidence and cost of nurse workplace violence perpetrated by hospital patients or patient visitors. *J Emerg Nurs*. 2014 May;40(3):218–28. <http://dx.doi.org/10.1016/j.jen.2013.05.014> PMID:24054728
- (21) Muzembo BA, Mbutshu LH, Ngatu NR, Malonga KF, Eitoku M, Hirota R, et al. Workplace violence towards Congolese health care workers: a survey of 436 healthcare facilities in Katanga province, Democratic Republic of Congo. *J Occup Health*. 2015;57(1):69–80.  
<http://dx.doi.org/10.1539/joh.14-0111-OA> PMID:25476862
- (22) Alsaleem SA, Alsabaani A, Alamri RS, Hadi RA, Alkhayri MH, Badawi KK, et al. Violence towards healthcare workers: A study conducted in Abha City, Saudi Arabia. *JFCM*. 2018 Sep-Dec;25(3):188–93. [http://dx.doi.org/10.4103/jfcm.JFCM\\_170\\_17](http://dx.doi.org/10.4103/jfcm.JFCM_170_17) PMID:30220849
- (23) Vezyridis P, Samoutis A, Mavrikiou PM. Workplace violence against clinicians in Cypriot emergency departments: a national questionnaire survey. *J Clin Nurs*. 2015 May;24(9-10):1210–22. <http://dx.doi.org/10.1111/jocn.12660> PMID:25047646
- (24) Cheung T, Lee PH and Yip PS. Workplace violence toward physicians and nurses: prevalence and correlates in Macau. *Int J Environ Res Public Health*. 2017 Aug 4;14(8). pii: E879. <http://dx.doi.org/10.3390/ijerph14080879> PMID:28777333
- (25) Cheung T and Yip PS. Workplace violence towards nurses in Hong Kong: prevalence and correlates. *BMC Public Health*. 2017 Feb 14;17(1):196. <http://dx.doi.org/10.1186/s12889-017-4112-3> PMID:28196499
- (26) Abbas MA, Fiala LA, Abdel Rahman AG and Fahim AE. Epidemiology of workplace violence against nursing staff in Ismailia Governorate, Egypt. *J Egypt Public Health Assoc*. 2010 Jan 1;85(1–2):29–43. PMID:21073846
- (27) Jmal-Hammami K, Loukil-Feki M, Moalla E, Gargouri I, Masmoudi ML, Marouen-Jamoussi S. Les agressions sur les lieux du travail en milieu hospitalier: À propos de 107 cas. *Archives des Maladies Professionnelles et de l'Environnement*. 2006 Sep 1;67(4):626–30.  
[https://doi.org/10.1016/S1775-8785\(06\)70441-6](https://doi.org/10.1016/S1775-8785(06)70441-6)
- (28) Gillespie GL, Gates DM, Miller M and Howard PK. Workplace violence in healthcare settings: risk factors and protective strategies. *Rehabilitation nursing*. 2010 Sep–Oct;35(5):177–84. <http://dx.doi.org/10.1002/j.2048-7940.2010.tb00045.x> PMID:20836482
- (29) Hogarth KM, Beattie J and Morphet J. Nurses' attitudes towards the reporting of violence in the emergency department. *Australas Emerg Nurs J*. 2016 May;19(2):75–81.  
<http://dx.doi.org/10.1016/j.aenj.2015.03.006> PMID:26012889

**Table 1. Demographic and occupational characteristics of HCWs in EDs, Dammam**

HCW characteristics	No.	%
Occupation		
Physician	63	19
Nurse	175	54
Others <sup>a</sup>	86	27
Sex		
Male	109	33.6
Female	215	66.4
Age (yr) <sup>b</sup>		
≤ 30	153	47.2
31–40	136	42
> 40	35	10.8
Marital status		
Married	226	69.8
Unmarried	98	30.2
Nationality		
Saudi	253	78.1
Non-Saudi	71	21.9
Education		
Diploma	163	50.3
Bachelor's	141	43.5
Master's	12	3.7
Board <sup>c</sup>	8	2.5
Work experience (yr)		
1–5	126	38.9
6–9	130	40.1
> 10	68	21
Multiple shifts		
Yes	292	90.1
No	32	9.9
Shift time		
Morning	42	13
Alternate	282	87
No. of coworkers		
Mean (standard deviation)	9 (5)	
≤ 10	232	71.6
> 10	92	28.4

<sup>a</sup>Pharmacists, technicians and clerical workers.

<sup>b</sup>Mean age 32.7 (6.2) years.

<sup>c</sup>Medical degree for physicians to receive privileges and to practice medicine in a particular field.

ED = emergency department; HCW = healthcare worker.

**Table 2. Characteristics and types of workplace violence among HCWs in EDs, Dammam**

Characteristics and types of violence	Physical		Verbal		Bullying		Sexual		Racial		Total	
	n	%	n	%	n	%	n	%	n	%	n	%
	45	19	126	52	39	16	7	3	24	10	241	100
Shift time												
Morning	10	22.2	42	33.3	29	74.4	3	42.9	11	45.8	95	39.4
Evening	22	48.9	55	43.7	6	15.4	3	42.9	9	37.5	95	39.4
Night	13	28.9	29	23	4	10.3	1	14.3	4	16.7	51	21.2
Location												
Inside	40	88.9	123	97.6	38	97.4	7	100	24	100	232	96.3
Outside	0	0	1	0.8	1	2.6	0	0	0	0	2	0.8
Both	5	11.1	2	1.6	0	0	0	0	0	0	7	2.9
Frequency <sup>a</sup>												
Once a year	24	53.3	39	31	21	53.8	5	71.4	10	41.7	99	41.1
Once a month	14	31.1	39	31	9	23.1	0	0	7	29.2	69	28.6
More than once per month	7	15.6	48	38	9	23.1	2	28.6	7	29.2	73	30.3
Offender identity												
Patient / client	24	53.3	65	51.6	0	0	2	28.6	11	45.8	102	42
Relatives	16	35.6	48	38.1	0	0	1	14.3	10	41.7	75	31.1
Staff member <sup>b</sup>	1	2.2	5	4	14	35.9	3	42.9	0	0	23	10
Management	0	0	1	0.8	22	56.4	0	0	3	12.5	26	10.8
External colleague	1	2.2	1	0.8	3	7.7	0	0	0	0	5	2
General public	3	6.7	6	4.8	0	0	1	14.3	0	0	10	4.1
Offender age												
< 20 years	0	0	3	2.4	0	0	0	0	0	0	3	1
21–45 years	42	93.3	100	79.4	30	76.9	7	100	18	75	197	82
≥ 46 years	3	6.7	23	18.3	9	23.1	0	0	6	25	41	17
Offender sex <sup>a</sup>												
Male	23	51.1	40	32	18	46.2	5	71.4	9	37.5	95	39.42
Female	7	15.6	24	19	11	28.2	2	28.6	5	20.5	49	20.33
Both	15	33.3	62	49	10	25.6	0	0	10	41.7	97	40.25
Could have been prevented												
Yes	41	91.1	91	72.2	28	71.8	4	57.1	16	66.7	180	74.7
No	4	8.9	35	27.8	11	28.2	3	42.9	8	33.3	61	25.3
Consequences on attacker												
None	19	42.2	80	63.5	34	87.2	4	57.1	17	70.8	154	63.9
Verbal warning	13	28.9	29	23	2	5.1	3	42.9	4	16.7	51	21.2
Reported to police	9	20	7	5.6	0	0	0	0	0	0	16	6.6
Do not know	4	8.9	10	7.9	3	7.7	0	0	3	12.5	20	8.3
Consequences on HCW												
None	16	35.6	61	48.4	20	51.3	5	71.4	10	41.7	112	46.5
Reduce work performance	20	44.4	54	42.9	18	46.2	2	28.6	13	54.2	107	44.4
Incident form against HCWs	7	15.6	11	8.7	1	2.6	0	0	1	4.2	20	8.3
Injured	2	4.4	0	0	0	0	0	0	0	0	2	0.8

<sup>a</sup>Percentage calculated from HCWs that could have been subjected to > 1 incidence of violence.

<sup>b</sup>Physicians, nurses, pharmacists, technicians and clerical workers.

ED = emergency department; HCW = healthcare worker.

**Table 3. History of workplace violence related to characteristics of HCWs in EDs, Dammam**

HCW characteristics	History of exposure to violence				Total 324		$\chi^2$	P
	Yes		No					
	n	%	n	%	n	%		
Sex								
Male	63	57.8	46	42.2	109	33.6	6.528	0.01
Female	92	42.8	123	57.2	215	66.4		
Age	155		169		324			0.622
Mean (SD)	32.5 (5.7)		32.9 (6.6)					
Age group, yr								
≤ 30	70	45.8	83	54.2	153	47.2	2.168	0.338
31–40	71	52.2	65	47.8	136	42		
> 40	14	40	21	60	35	10.8		
Marital status								
Married	109	48.2	117	51.8	226	69.8	0.046	0.831
Unmarried	46	46.9	52	53.1	98	30.2		
Nationality								
Saudi	131	51.8	122	48.2	253	78.1	7.179	0.007
Non-Saudi	24	33.8	47	66.2	71	21.9		
Occupation								
Physician	34	54	29	46	63	19	1.75	0.417
Nurse	84	48	91	52	175	54		
Others	37	43	49	57	86	27		
Shift time								
Morning	17	40.5	25	59.5	42	13	1.048	0.306
Alternate	138	48.9	144	51.1	282	87		
No. of coworkers								
≤ 10	124	53.4	108	46.6	232	71.6	10.3	0.001
> 10	31	33.7	61	66.3	92	28.4		
Report encouragement								
Yes	76	39.4	117	60.6	193	59.6	13.69	<0.001
No	79	60.3	52	39.7	131	40.4		
System availability								
Yes	98	43.6	127	56.4	225	69.4	5.41	0.02
No	57	57.6	42	42.4	99	30.6		

ED = emergency department; HCW = healthcare worker; SD = standard deviation.

**Table 4. Type of workplace violence related to characteristics of HCWs in EDs, Dammam**

HCW characteristics	Physical						
	Yes	%	No	%	Total 324	$\chi^2$	<i>P</i>
Sex							
Male	22	20.2	87	79.8	109	5.442	0.02
Female	23	10.7	192	89.3	215		
No. of coworkers							
≤ 10	38	16.4	194	83.6	232	4.237	0.04
> 10	7	7.6	85	92.4	92		
	Verbal						
Sex							
Male	51	46.8	58	53.2	109	4.314	0.038
Female	75	34.9	140	65.1	215		
Nationality							
Saudi	106	41.9	147	58.1	253	4.397	0.036
Non-Saudi	20	28.2	51	71.8	71		
No. of coworkers							
≤ 10	101	43.5	131	56.5	232	7.420	0.006
> 10	25	27.2	67	72.8	92		
Report encouragement							
Yes	61	31.6	132	68.4	193	10.653	0.001
No	65	49.6	66	50.4	131		
	Bullying						
Report encouragement							
Yes	16	8.3	177	91.7	193	6.329	0.012
No	23	17.6	108	82.4	131		
	Sexual						
Marital status							
Married	2	0.9	224	99.1	226		0.028
Unmarried	5	5.1	93	94.9	98		

ED = emergency department; HCW = healthcare worker; SD = standard deviation.

**Table 5. Logistic regression analysis of workplace violence using significantly associated characteristics of HCWs in EDs, Dammam**

Variables		B	SE	Wald test	df	Sig	Exp(B)	95% CI for Exp(B)	
								Lower	Upper
General	Saudi nationality	1.015	0.438	5.375	1	0.020	2.759	1.170	6.507
	Lack of report	-0.915	0.375	5.945	1	0.015	2.497	1.197	5.209
	encouragement	-0.351	0.723	0.236	1	0.627	0.704		
	Constant								
Physical	Male sex	1.045	0.485	4.632	1	0.031	2.842	1.098	7.358
	Constant	-2.380	0.331	51.824	1	< 0.001	0.093		
Verbal	Lack of report	0.887	0.425	4.350	1	0.037	2.428	1.055	5.589
	encouragement	-2.260	0.595	14.448	1	<0.001	0.104		
	Constant								
Bullying	Lack of report	0.857	0.348	6.074	1	0.014	2.356	1.192	4.657
	encouragement	-3.260	0.570	32.676	1	< 0.001	0.038		
	Constant								

*df* = degrees of freedom; *ED* = emergency department; *HCW* = healthcare worker; *SE* = standard error; *Sig* = significance.