

Table 4 Logistic regression models of diabetes and hypertension according to waist circumference (WC), waist-to-hip ratio (WHR) and body mass index (BMI) among Egyptian adults

Obesity indicator/ Morbidity	β (SE)	<i>P</i>-value	Exp. β	95% CI
<i>Waist circumference</i>				
Diabetes mellitus ^a	0.02 (0.01)	0.02	1.02	1.01–1.04
Hypertension ^b	0.02 (0.01)	0.0007	1.02	1.01–1.04
<i>Waist-to-hip ratio</i>				
Diabetes mellitus ^c	-1.49 (1.64)	0.36	0.23	0.01–5.57
Hypertension ^d	-0.39 (1.02)	0.70	0.68	0.09–4.98
<i>Body mass index</i>				
Diabetes mellitus ^e	0.01 (0.01)	0.11	1.01	0.99–1.02
Hypertension ^f	-0.01 (0.01)	0.96	0.99	0.98–1.02

^aAdjusted for age, sex, hypertension and smoking; ^bAdjusted for age, sex and diabetes, smoking and WHR; ^cAdjusted for age, sex, hypertension, smoking and WC; ^dAdjusted for age, sex, diabetes, smoking and WC; ^eAdjusted for age, sex, hypertension, smoking WC and WHR; ^fAdjusted for age, sex and diabetes, smoking, WC and WHR.

SE = standard error; CI = confidence interval.