interactive group discussions			
Item	Pre-intervention	Post-intervention	
	All physicians	Control group	Intervention group

Mean (SD) score

6.00 (1.00)

0.46 (0.09)

0.86 (0.12)

0.67 (0.11)

0.99 (0.13)

0.95 (0.08)

0.67 (0.11)

0.68 (0.09)

0.86 (0.09)

Mean (SD) score

6.24 (0.80)

0.52 (0.11)

0.85 (0.11)

0.69(0.14)

0.99(0.10)

0.98 (0.11)

0.69 (0.12)

0.71 (0.11)

0.88 (0.11)

Table 1 Scores for correct principles of prescription writing attained by all physicians pre-intervention, postintervention in the control group, and post-intervention in the intervention group who participated in the

0.45 (0.11)

0.86(0.09)

0.67 (0.14)

0.97 (0.10)

0.95 (0.09)

0.68 (0.12)

0.66 (0.11)

0.85 (0.12)

	All physicians	
	Mean (SD) score	
Total prescribing pattern <sup>a</sup>	6.00 (1.10)	

Names of drugs written correctly and

Dosage forms of drugs written<sup>b</sup>

Directions for use written correctly<sup>b</sup>

<sup>a</sup>Maximum score 8: <sup>b</sup>maximum score 1.

Directions for use written completely<sup>b</sup>

Strength of drugs written<sup>b</sup>

Number of drugs written<sup>b</sup>

Directions for use writtenb

All items written clearly<sup>b</sup>

SD = standard deviation.

completely<sup>b</sup>