

Table 3 Clinical, echocardiographic and laboratory data for dialysis patients with and without ST-segment depression

Parameter	ST-depression (n = 28)		No ST-depression (n = 20)	
	Mean	SD	Mean	SD
Age (years)	30.1	2.2	30.1	3.1
Serum Na ⁺ (mmol/L)				
Before dialysis	136.2	2.5	135.7	2.6
After dialysis	135.9	2.2	136.7	2.0
Serum K ⁺ (mmol/L)				
Before dialysis	5.3	0.4	5.3	0.3
After dialysis	3.6	0.9	4.5	1.1**
Serum Ca ²⁺ (mg/dL)				
Before dialysis	1.97	0.12	2.00	0.17
After dialysis	2.59	0.22	2.57	0.22
Serum Mg ²⁺ (mg/dL)				
Before dialysis	1.46	0.09	1.41	0.07
After dialysis	1.45	0.07	1.42	0.06
LVM (g)	244.2	101.9	177.4***	44.2
LVMI (g/m ²)	162.6	90.9	88.6*	28.3
Male sex [No. (%)]	20 (71%)		12 (60%)	
Hypertension [No. (%)]	15 (54%)		4 (20%)*	

*P-value < 0.05; **P-value < 0.01; ***P-value < 0.001.

SD = standard deviation.

LVM = left ventricular mass; LVMI = left ventricular mass index.