Table 6 Prevalence of tobacco use in the hosting countries and Eastern Mediterranean Region and smoking detection rates in the present study of Palestine refugees

| 0 | | | |
|---|-------|---------|------------------------------------|
| Country/population | Males | Females | Source |
| | % | % | |
| Jordan | | | |
| Host country | 50.5 | 8.3 | 2002 national survey, age > 18 yr |
| Palestine refugees ^a | 48.0 | 8.4 | UNRWA 2007 |
| Lebanon | | | |
| Host country | 60.7 | 46.9 | 1997 national survey, age > 19 yrs |
| Palestine refugees ^a | 59.5 | 39.9 | UNRWA 2007 |
| Gaza Strip and West Bank | | | |
| Host country | 20.0 | 18.0 | EMRO country profile 2004 |
| Gaza: Palestine refugees* | 49.5 | 0.6 | UNRWA 2007 |
| West Bank: Palestine refugees ^a | 43.7 | 6.1 | UNRWA 2007 |

21.0

2004 Aleppo survey, age 18-65 vrs

Palestine refugees^a 57.1 15.3 **UNRWA 2007** Israel 38.6 22.1 2001 national survey, age 25-64 yrs Eastern Mediterranean Region 34.2 8.7 1998 WHO UNRWA = United Nations Relief and Works Agency for Palestine Refugees in the Near East; EMRO = World Health Organization Regional Office for the Eastern Mediterranean; WHO - World Health Organization. *Data refer only to the Palestinian refugees screened for noncommunicable diseases at UNRWA health centres.

62.0

Syrian Arab Republic Host country (subnational)