

**Table 3 Coverage of malaria preventive and curative services as reflected by the findings of the demographic and health survey of Pakistan 2006-07**

Variable/indicator	Urban <i>n</i> (%)	Rural <i>n</i> (%)	Total <i>n</i> (%)
Household possession of nets (treated or untreated)	3159 (3.5)	6096 (7.8)	9255 (6.3)
Households with at least one insecticide-treated net (ITN)	3159 (0.7)	6096 (0.8)	9255 (0.8)
Children under 5 years who slept under a mosquito net the night before the interview	2636 (1.2)	6142 (1.8)	8778 (1.6)
Children under 5 years who slept under an ITN the night before the interview	2636 (0.2)	6142 (0.2)	8778 (0.2)
Pregnant women who slept under a mosquito net the night before the interview	364 (0.7)	829 (2.0)	1193 (1.6)
Pregnant women who slept under an ITN the night before the interview	364 (0.0)	829 (0.0)	1193 (0.0)
Children under 5 years with fever in the two weeks preceding the survey who took antimalarial drugs	791 (2.5)	1777 (3.7)	2569 (3.3)
Children under 5 years with fever in the two weeks preceding the survey who took antimalarial drugs the same day/next day	791 (2.0)	1777 (2.8)	2569 (2.6)