

Table 1 Relationship between some sociodemographic characteristics and presence of erectile dysfunction among men aged > 30 years attending primary care centres in Qatar for any reason

Variable	Erectile dysfunction		Normal		χ^2	P-value
	No.	%	No.	%		
Age (years)						
30-40	153	40.7	223	59.3	101.33	< 0.001
> 40-50	254	57.5	188	42.5		
> 50-60	179	71.9	70	28.1		
> 60	63	88.7	8	11.3		
Nationality						
Qatari	110	64.7	60	35.3	9.84	< 0.001
Other Arab	183	48.4	195	51.6		
Asian	349	60.7	226	39.3		
Other	7	43.8	9	56.3		
Education						
Illiterate	174	64.7	95	35.3	8.72	0.012
Pre-university	280	55.4	225	44.6		
University and above	195	53.4	170	46.6		
Occupation						
Unemployed	62	79.5	16	20.5	78.17	< 0.001
Clerk	216	68.8	98	31.2		
Army	12	34.3	23	65.7		
Skilled manual	319	48.0	345	52.0		
Retired	40	83.3	8	16.7		
Income					7.27	0.026
Satisfactory	442	59.6	299	40.4		
Unsatisfactory	207	52.0	191	48.0		
No. of children					19.66	< 0.001
0	51	7.9	51	10.4		
1-3	302	46.5	279	56.9		
> 3	296	45.6	160	32.7		