

Table 2 Trends in high risk sexual and injecting behaviours among most at-risk populations in Pakistan

Variable	Female sex workers		Male sex workers			Hijra sex workers			Injection drug users		
	2005-06	2006-07	2005-06	2006-07	2008	2005-06	2006-07	2008	2005-06	2006-07	2008
No of cities	8	12	8	12	8	8	12	8	8	12	9
No of interviews	3134	4639	1781	2289	1205	1569	2143	1186	2432	4039	2979
Mean (SD) no. of clients/day	4.5	2.6 (1.8)	2.3	2.1 (1.5)	1.9 (0.9)	2.5	2.5 (2.1)	2.6 (1.9)	-	-	-
Consistent condom use (%)	18.0	23.0	6.8	7.0	24.0	5.8	7.5	19.7	-	-	-
Condom use at last commercial sexual encounter (%)	34.0	45.0	24.0	21.0	35.0	21.0	19.0	32.0	17.0	21.0	31.0
Sex with injection drug users (%)	13.5	9.9	10.1	9.5	6.4	8.4	6.4	6.3	-	-	-
Shared needle at last injection (%)	-	-	-	-	-	-	-	-	47.1	21.9	17.6
Injected with used needle (%)	-	-	-	-	-	-	-	-	35.2	27.8	22.5
Injection drug users having sex with female sex workers in past 6 months (%)	-	-	-	-	-	-	-	-	12.6	26.6	17.7
Coverage of prevention programmes (%)	2.1	1.9	2.2	2.7	8.5	2.5	4.2	18.3	19.1	15.6	50.8

-- data not applicable.
SD = standard deviation.