

Table 4 Mean scores of participants on the scopes and indices of the SCL-90-R test ($n = 687$)

Scopes and indices of SCL-90-R Test	Total Patients Mean (SD) scores ($n = 687$)	Group A Mean (SD) scores ($n = 405$)	Group B Mean (SD) scores ($n = 282$)
Somatization	0.97 (0.61)	0.64 (0.37)	1.43 (0.59)
Obsessive-compulsive	0.96 (0.65)	0.59 (0.38)	1.48 (0.60)
Interpersonal sensitivity	1.12 (0.72)	0.7 (0.41)	1.74 (0.64)
Depression	1.13 (0.76)	0.66 (0.38)	1.8 (0.67)
Anxiety	0.9 (0.66)	0.53 (0.35)	1.42 (0.64)
Hostility	0.96 (0.74)	0.59 (0.45)	1.48 (0.76)
Phobic anxiety	0.64 (0.58)	0.36 (0.32)	1.04 (0.63)
Paranoid ideation	1.26 (0.78)	0.84 (0.51)	1.86 (0.71)
Psychoticism	0.78 (0.61)	0.43 (0.33)	1.28 (0.57)
Global severity index	0.97 (0.02)	0.6 (0.25)	1.51 (0.44)
Positive symptom total	47.64 (0.73)	36.25 (14.37)	64 (12.18)
Positive symptom distress index	1.77 (0.01)	1.52 (0.38)	2.12 (0.47)

Group A: probable safe subjects having lower global severity index than the mean score of the sample in this research

Group B: probable patient subjects having higher global severity index than the mean score of the sample in this research.

SD = standard deviation.