Research intervention All participants Sex **Educational level** Income (JD) (n = 120)Men Women School College/ < 500 ≥ 500

(n = 73)

(n = 22)

42.1**

university

16.7

(n = 44)

31.6*

(n = 76)

15.5

Table 2 Respondents' preferences for participating in research involving different types of interventions: percentage of respondents

(n = 47)

31.7**

21.1

who maybe or certainly would participate

Surgical procedures

*P < 0.05; **P < 0.01. JD = Jordanian dinars.

| | | | | | (n = 98) | | |
|---------|------|------|------|------|----------|------|------|
| | % | % | % | % | % | % | % |
| Surveys | 81.7 | 83.7 | 80.3 | 72.2 | 83.5 | 82.1 | 81.4 |

| Surveys | 81./ | 83./ | 80.3 | /2.2 | 83.5 | 82.1 | 81.4 |
|-------------------|------|------|------|------|------|------|------|
| Oral swabs | 52.8 | 57.5 | 50.0 | 50.0 | 53.3 | 50.0 | 54.3 |
| Saliva collection | 51.4 | 56.1 | 48.5 | 52.6 | 51.1 | 44.7 | 54.9 |

| Orar swabs | 32.0 | 37.3 | 50.0 | 50.0 | 33.3 | 50.0 | 57.5 | |
|---------------------|------|------|------|------|------|------|------|--|
| Saliva collection | 51.4 | 56.1 | 48.5 | 52.6 | 51.1 | 44.7 | 54.9 | |
| Topical medications | 50.5 | 56.1 | 47.1 | 52.6 | 50.0 | 60.5 | 45.1 | |
| | | | | | | | | |

| Topical medications | 50.5 | 56.1 | 47.1 | 52.6 | 50.0 | 60.5 | 45.1 | |
|---------------------|------|------|------|------|------|------|------|--|
| Restoration | 44.4 | 53.7 | 38.8 | 57.9 | 41.6 | 50.0 | 41.4 | |
| | | | | | | | | |

| Restoration | 44.4 | 53./ | 38.8 | 57.9 | 41.6 | 50.0 | 41.4 |
|-------------|------|------|------|------|------|------|------|
| Biopsies | 22.0 | 24.4 | 20.6 | 21.1 | 22.2 | 18.4 | 23.9 |

14.7