Table 3 Clinical presentation in the case and control groups before and after treatment												
Symptom	Case group ( <i>n</i> = 144)					Control group ( <i>n</i> = 140)					Statistical analysis	
	Before		After		<i>P</i> -value <sup>a</sup>	Before		After		<i>P</i> -value <sup>a</sup>	RR (95% CI)	<i>P</i> -value <sup>b</sup>
	No.	%	No.	%		No.	%	No.	%			
Cough	127	88	101	70	0.004	112	80	103	74	0.427	2.3 (1.3-4.1)	0.003
Wheezing	94	65	68	47	0.002	88	63	81	58	0.638	3.6 (1.8–7.4)	< 0.001
Dyspnoea	56	39	35	24	0.003	57	41	49	35	0.537	4.2 (1.9-8.7)	< 0.001
Partially controlled asthma <sup>c</sup>	144	100	73	51	< 0.001	140	100	112	80	0.261	2.0 (1.4-2.8)	< 0.001

<sup>*a*</sup>Before versus after,  $\chi^2$ -test; <sup>*b*</sup>Case versus control group, independent t-test.

<sup>c</sup>Daytime symptoms more than twice/week; limitations of activities and nocturnal symptoms/activities; need for relief/rescue therapy more than twice/week; lung function (PEF or FEV1) < 80% of predicted or personal best. RR = relative risk; 95% CI = confidence interval.