

Table 1 Characteristics of hospitalized patients with HIV/AIDS

Variable	Total	Male		Female		P-value
<i>Total admissions (No., %)</i>	227	100.0	187	82.4	40	17.6
<i>Age (years) [Median, (IQR)]</i>	40	(36–45)	40	(37.5–45)	36.5	(30–45)
<i>Marital status (No., %)</i>						< 0.001 <sup>c</sup>
Single	110	48.5	104	55.6	6	15.0
Married	84	37.0	61	32.6	23	57.5
Divorced	10	4.4	9	4.8	1	2.5
Widowed	9	4.0	0	0.0	9	22.5
Unknown	14	6.2	13	7.0	1	2.5
<i>New HIV diagnosis (No., %)</i>	32	14.1	23	12.3	9	22.5
<i>HIV risk factors (No., %)</i>						< 0.001 <sup>c</sup>
Injection drug use	109	48.0	109	58.3	0	0.0
Heterosexual	40	17.6	5	2.7	35	87.5
Homosexual	2	0.9	2	1.1	0	0.0
Unknown	76	33.5	71	38	5	12.5
<i>HAART use (No., %)</i>						0.8 <sup>c</sup>
Never	103	45.4	83	44.4	20	50.0
Current	82	36.1	69	36.9	13	32.5
Defaulted	38	16.7	32	17.1	6	15.0
Unknown	4	1.8	3	1.6	1	2.5
<i>Duration of hospital stay (days) [Median, (IQR)]</i>	8	(3–16)	9	(3–16)	7	(3–16.2)
<i>HCV-Ab positive (No., %)</i>	123	54.2	123	65.8	0	0.0
<i>HbsAg positive (No., %)</i>	11	4.8	11	5.9	0	0.0
<i>Hospital mortality (No., %)</i>	85	37.4	76	40.6	9	22.5
<i>Death within 48 h of admission (No., %)</i>	30	13.2	28	15.0	2	5.0

<sup>a</sup>Mann-Whitney U test; <sup>b</sup>Chi-squared-test; <sup>c</sup>Fisher exact test.

IQR = interquartile range; HAART = highly active antiretroviral therapy; HCV-Ab = anti-hepatitis C virus antibody; HBsAg = hepatitis B virus surface antigen.