

Table 3 Association of awareness of folic acid with selected demographic and obstetric characteristics

Variable	Bivariate analysis		Multivariable analysis	
	OR (95%CI)	P-value	OR (95%CI)	P-value
Age (years)				
< 18*				
18–	1.24 (0.27–5.62)	0.779	0.45 (0.08–2.44)	0.35
31–	1.55 (0.34–7.12)	0.572	0.85 (0.41–1.77)	0.66
≥ 41	2.29 (0.46–11.36)	0.312	0.97 (0.47–1.99)	0.93
Residence				
Rural*				
Urban	2.17 (1.57–2.99)	< 0.001	2.17 (1.51–3.21)	< 0.001
Education				
Illiterate*				
Primary	1.097 (0.71–1.7)	0.676	1.9 (0.89–4.08)	0.096
Secondary	0.667 (0.42–1.05)	0.082	1.8 (0.87–3.84)	0.113
Higher secondary	0.656 (0.38–1.12)	0.125	1.1 (0.54–2.44)	0.725
University and above	0.45 (0.22–0.94)	0.032	1.1 (0.49–2.48)	0.806
Occupation				
Employee*				
Housewife	0.32 (0.14–0.75)	0.009	1.61 (0.20–13.22)	0.660
Labourer	0.83 (0.34–2.04)	0.688	0.71 (0.07–6.85)	0.768
Self-employed business woman	0.45 (0.06–0.55)	0.446	1.23 (0.12–12.40)	0.862
Parity				
1*				
1–	0.79 (0.50–1.25)	0.306	0.88 (0.53–1.45)	0.609
≥ 5	1.51 (1.01–2.28)	0.047	0.63 (0.34–1.17)	0.142
Prior abortion				
Yes*				
No	1.02 (0.71–1.46)	0.902	1.01(0.69–0.49)	0.951
Antenatal care (at least one visit)				
No*				
Yes	2.30 (1.46–3.78)	< 0.001	2.20 (1.27–3.96)	0.005
Antenatal care entry				
Not used*				
1st trimester	0.44 (0.27–0.73)	0.001	0.58 (33–1.02)	0.059
2nd trimester	1.16 (0.67–2.00)	0.596	1.35 (0.73–2.49)	0.341
3rd trimester	1.32 (0.70–2.47)	0.392	1.75 (0.88–3.48)	0.114
Booked				
Yes*				
No	1.50 (1.04–2.08)	0.030	0.85 (0.56–1.29)	0.449

*Reference category.

OR = odd ratios; CI = confidence interval.