

Table 3 Adjusted ORs (95% confidence interval) for correlating factors for smoking contemplation and maintenance

	Contemplation		Maintenance	
	Cigarette	Waterpipe	Cigarette	Waterpipe
<i>Depression</i>	1.96 (1.41–2.72)	1.09 (0.89–1.33)	1.49 (0.83–2.68)	0.87 (0.58–1.31)
<i>Risky behaviour</i>	2.13 (1.51–2.94)	1.49 (1.22–1.85)	3.70 (1.85–7.14)	1.89 (1.33–3.03)
<i>Family conflict</i>	1.87 (1.38–2.53)	1.53 (1.24–1.88)	1.10 (0.59–2.05)	1.25 (0.83–1.89)
<i>More positive attitude to smoking acceptability</i>	2.12 (1.51–2.97)	1.72 (1.41–2.10)	1.46 (0.73–2.91)	1.57 (1.03–2.40)
<i>Self-efficacy</i>				
Low	1	1	1	1
Medium	0.75 (0.41–0.80)	0.78 (0.61–0.96)	1.46 (0.78–2.73)	1.47 (0.95–2.26)
High	0.47 (0.29–0.85)	0.79 (0.94–1.70)	0.86 (0.39–1.93)	1.14 (0.62–2.09)
<i>Log likelihood</i>	1297.065	2713.739	1324.466	2794.327
<i>R²</i>	0.35	0.38	0.36	0.38

Odds ratios adjusted for age, sex, parental education and smoking.

Bold numbers with $P < 0.05$.