

Table 2. Association between conflict exposure and quality-of-life domains (European Organization for Research and Treatment of Cancer Quality-of-life Questionnaire-Core 30)

Quality-of-life	Sample size	Mean score (SD) for low conflict exposure	Mean score (SD) for high conflict exposure	P value
Physical functioning	21	70.2 (18.6)	52.1 (23.4)	< 0.01
Role functioning	20	65.7 (20.1)	45.2 (25.6)	< 0.01
Emotional functioning	19	55.3 (22.4)	36.1 (20.3)	< 0.01
Cognitive functioning	37	60.2 (19.8)	46.9 (21.2)	< 0.01
Social functioning	14	63.4 (21.0)	40.8 (22.1)	< 0.01
Pain	26	28.7 (22.5)	47.8 (26.3)	< 0.01
Fatigue	25	54.5 (20.7)	72.1 (23.8)	< 0.01
Nausea/vomiting	115	18.2 (14.7)	23.9 (16.2)	0.05
Dyspnoea	57	14.9 (13.8)	23.2 (17.6)	0.01
Insomnia	32	38.6 (20.2)	53.8 (23.4)	< 0.01
Appetite loss	70	28.1 (22.0)	39.4 (25.6)	< 0.01
Financial difficulties	42	35.4 (26.2)	52.7 (30.1)	< 0.01

Results from t tests comparing quality-of-life scores between groups with low and high conflict exposure.