

Table 1 Characteristics of women diagnosed with invasive cervical cancer in Saudi Arabia, 2005–2019 (N = 2496)

		Saudi								Non-Saudi							
		2005–2009		2010–2014		2015–2019		All periods		2005–2009		2010–2014		2015–2019		All periods	
		mean	SD	mean	SD	mean	SD	mean	SD	mean	SD	mean	SD	mean	SD	mean	SD
Age		52.2	13.9	52.8	12.9	54.8	13.7	53.4	13.5	47.2	12.5	49.6	12.0	49.4	12.3	48.8	12.3
		n	%	n	%	n	%	n	%	n	%	n	%	n	%	n	%
Region of diagnosis	Riyadh	123	28.0	136	25.0	188	30.2	447	27.8	76	28.8	70	20.8	88	30.6	234	26.3
	Makkah	132	30.0	160	29.4	193	31.0	485	30.2	147	55.7	181	53.7	143	49.7	471	53.0
	Eastern	76	17.3	84	15.4	122	19.6	282	17.6	16	6.1	45	13.4	29	10.1	90	10.1
	Other	109	24.8	165	30.3	119	19.1	393	24.5	25	9.5	41	12.2	28	9.7	94	10.6
Stage	Localised	125	28.4	151	27.7	258	41.5	534	33.2	85	32.2	155	46.0	156	54.2	396	44.5
	Regional	198	45.0	220	40.4	191	30.7	609	37.9	70	26.5	73	21.7	74	25.7	217	24.4
	Distant	76	17.3	101	18.5	128	20.6	305	19.0	28	10.6	33	9.8	33	11.5	94	10.6
	Unknown	41	9.3	73	13.4	45	7.2	159	9.9	81	30.7	76	22.6	25	8.7	182	20.5
Grade	I	30	6.8	50	9.2	52	8.4	132	8.2	18	6.8	20	5.9	31	10.8	69	7.8
	II	159	36.1	172	31.6	185	29.7	516	32.1	76	28.8	126	37.4	104	36.1	306	34.4
	III	112	25.5	159	29.2	196	31.5	467	29.1	78	29.6	88	26.1	77	26.7	243	27.3
	IV	19	4.3	11	2.0	15	2.4	45	2.8	14	5.3	21	6.2	13	4.5	48	5.4
	Unknown	120	27.3	153	28.1	174	28.0	447	27.8	78	29.6	82	24.3	63	21.9	223	25.1
Basis of diagnosis	Clinical/ imaging	10	2.3	9	1.7	34	5.5	53	3.3	10	3.8	7	2.1	5	1.7	22	2.5
	Pathology	426	96.82	530	97.25	580	93.3	1536	95.6	252	95.5	328	97.3	281	97.6	861	96.9
	Death certificate only	4	0.9	6	1.1	8	1.3	18	1.1	2	0.8	2	0.6	2	0.7	6	0.7
Morphology	Squamous	304	69.1	362	66.4	397	63.8	1063	66.2	187	70.83	228	67.7	202	70.1	617	69.4
	Glandular	90	20.5	96	17.6	139	22.4	325	20.2	39	14.8	61	18.1	52	18.1	152	17.1
	Other	46	10.5	87	16.0	86	13.8	219	13.6	38	14.4	48	14.2	34	11.8	120	13.5