Canada, 2006 [52] Refugees 68 0.0 Planned Denmark, 1995 [53] Refugees 1936 6.71 Planned Holland, 1994-197 [54] Asylum seekers 46 424 2.22 Routine Holland, 1998-02 [30] Regular immigrants 68 122 1.16 Routine 6 Ireland, 1999-2000 [55] Refugees 945 12.7 12 Planned Kuwait, 1997-2006 [56] Regular immigrants 2 328 582 1.98 Routine 31 [57] Asylum seekers 900 4.44 New Zealand, 1999-2000 11 Routine Norway, 1987-95 [21] Asylum seekers 19912 2.16 8 Routine Norway, 1999 [58] Refugees 800 5.0 Planned Spain, 1998 [59] Regular immigrants 406 7.39 27 Planned

No. screened

1 863

4 794

TB prevalence among migrants

(cases per 1000 screened)

38.11

3.96

5.26

2.56

TB prevalence in host country

(cases per 100 000 populn)

6

15

6

Screening at entry

Planned

Planned

Planned

Planned

Routine

Routine

Routine

Routine

Routine

Planned

Planned

Planned

Switzerland, 1997-2004 [62] Regular immigrants 42 601 1.01 Switzerland, 2004 [63] Regular immigrants 8 9 9 5 5.56 UK, 2002-03 [64] Asylum seekers 8 258 1.33 12

Table 3 Patterns of screening for TB in migrants among selected host countries

Reference

[50]

[51]

[60]

[61]

Type of migrant

Screened refugees

Asylum seekers

Asylum seekers

Refugees

Host country, study date

Australia, 1999

Belgium, 1993

Switzerland, 2001-02

Switzerland, 1999

UK, 1995-99 [65] Asylum seekers 41 470 2.41 9 UK, 1998 [66] Regular immigrants 2 2 4 2 4.46 10 USA, 1980

13 507

3 1 1 9

<sup>[67]</sup> Regular immigrants 15 544 6.5 [68] Refugees 9 3 2 8 8.36

USA, 1980 19.5 USA, 1981-82 [69] Refugees 923

<sup>&</sup>quot;Routine" refers to standard screening protocol adopted by the local authorities; "Planned" refers to screening protocol specifically adopted to face a specific situation or for the purpose of a study. Source: Arshad et al. [28].