

Table 1 Distribution of surveyed women by sociodemographic and health service factors and percentages positive for indicators of antenatal care (ANC) utilization

Variable	All women (unweighted)		Adequate number of ANC visits ^{a,d}	Early use of ANC ^{b,d}	Skilled provider of ANC ^{c,d}
	No.	%	%	%	%
Woman's age (years)					
< 25	430	17.2	98.0	83.5***	98.3
25–34	1175	56.8	96.7	74.1	99.0
35+	547	26.0	96.3	70.7	99.6
Birth order					
1	468	21.6	98.7*	85.9***	98.8
2–4	868	41.3	97.5	75.2	98.4
5+	816	37.1	95.5	69.6	99.6
Place of residence					
Urban	1123	71.5	96.7	75.7	99.0
Rural	1029	28.5	97.0	72.6	99.3
Educational level					
Illiterate	350	9.5	94.6	67.7	97.0
Literate/primary/ preparatory	936	39.5	96.8	74.2	99.3
Secondary	635	34.7	97.4	77.5	99.0
University+	231	16.3	96.9	75.0	99.8
Economic status					
Low	373	15.8	96.2	76.6	98.0*
Lower middle	483	20.4	98.8	71.6	99.1
Upper middle	556	29.0	96.2	74.9	98.6
High	624	34.8	97.0	75.9	99.8
Work status^e					
Currently work	63	12.6	99.2	78.7	98.8
Previously worked	26	6.6	100.0	87.0	100.0
Never worked	503	80.8	97.5	74.8	99.5
Type of health facility^e					
Private	92	28.1	100.0	72.2	61.4
Public	293	71.9	97.0	76.3	70.7
Access time to health facility (min)^e					
≤ 30	307	88.2	97.5	74.4	67.1
> 30	61	11.8	100.0	79.5	68.4
Satisfaction with outpatient care					
Not satisfied	33	8.6	92.2	61.5	76.7
Satisfied	352	91.4	98.4	76.4	67.3
Total^f	2152	100.0	96.8	74.9	99.1

^aAdequate number of ANC visits – 4+ ANC visits; ^bEarly use of ANC – 1st antenatal visit in 1st trimester; ^cSkilled provider of ANC – care received from doctor or nurse/midwife.

^dRow percentages; ^eOnly those who attended health facilities during the previous 12 months; ^fTotals not always the same due to missing data in some variables.

* $P < 0.05$; *** $P < 0.001$.