

**Highlights**  
**WHO Eastern Mediterranean Region**

**Rapid assessment of service delivery for NCDs  
during the COVID-19 pandemic**

**Department of UHC/NCDs**  
**June 2020**



# Overview




- Rapid assessment sent to all Ministry of Health focal points who responded to the NCD country capacity survey in 2019
- Web-based questionnaire with 12 questions designed to assess
  - NCD staffing and service disruptions during the COVID-19 pandemic
  - services included in national COVID-19 response plans
  - funding for NCDs in national COVID-19 response plans
  - level of service disruption for a set of tracer services
  - main causes of disruption
  - mitigation strategies, data collection on co-morbidity
  - suggestions for WHO technical guidance and tools
- 86 % response rate (19 out of 22 countries in the Eastern Mediterranean)

# Survey questions




1. Are the Ministry of Health (or equivalent institutes) staff with responsibility for NCDs and their risk factors being reassigned/deployed to help with the overall COVID-19 response?
2. How much of the government (or Ministry of Health) funds initially allocated for NCDs has been reassigned to non-NCD services due to COVID-19 response efforts?
3. Is ensuring continuity of NCD services included in the list of essential health services in your country's COVID-19 response plan?
4. Which NCD services are included in the list of essential health services of your country's COVID-19 response plan? (CVD, cancer, diabetes, CRD, chronic kidney disease and dialysis, dental, rehab, tobacco-cessation, other)
5. Is there additional funding allocated for NCDs in the government budget for the COVID-19 response?
6. Which of the following Ministry of Health NCD activities planned for this year have been postponed because of COVID-19? (NCD surveys, public screening programmes, WHO PEN, HEARTS, mass communication campaigns, other)
7. During the COVID-19 pandemic, what are the government policies for access to essential NCD services at primary, secondary and tertiary care levels? (options assess outpatient and inpatient services)
8. Which of the following NCD-related services have been disrupted due to COVID-19? (HTN management, cardio emergencies, cancer Rx, diabetes management, asthma, urgent dental care, rehabilitation, palliative care)
9. What are the main causes of this disruption(s)? (options provided)
10. What approaches are being used to overcome the service disruptions to NCD management and prevention in public sector health facilities? (telemedicine, task shifting, novel supply chain or dispensing for meds, triaging, redirection to alternate services, other)
11. What are your country's plans to re-initiate any suspended NCD services?
12. Is the Ministry of Health collecting or collating data on NCD-related comorbidities in COVID-19 patients?
13. Are there any technical guidance or tools that you would suggest WHO to develop related to NCDs during the COVID-19 pandemic?


# NCD staff reassigned/deployed to help with COVID-19 response




# Funds for NCDs reassigned to non-NCD services due to COVID-19




## Continuity of NCD services included in essential services list in COVID-19 plan



# NCD services included in essential services list in COVID-19 plan




# Ministry of Health NCD planned activities postponed



# Access to essential NCD services at primary, secondary and tertiary levels




### Outpatient services




- Outpatient NCD services are closed
- Outpatient NCD services are open
- Outpatient NCD services are open with limited access and/or staff or in alternate locations or with different modes

### Inpatient services




- Inpatient NCD management services are open
- Inpatient NCD management services are open for emergencies only


# NCD-related services disrupted due to COVID-19




# Main causes of NCD service disruption



# Approaches to overcome service disruptions



# Data on NCD-related comorbidities in COVID-19 patients



## Suggested guidance or tools for WHO to develop on NCDs during COVID-19



- Inclusion of NCDs in all public health emergency protocols to ensure service continuity and to plan for service disruptions arising from long-term lockdowns
- Comprehensive toolkits for NCDs in emergencies (providers, patients, awareness, triage, tracing, linkages, referrals)
- Technical guidance on provision of ambulatory essential NCD services in case of lockdowns, without jeopardizing safety of patients or health care providers
- Health technology assessment of available digital solutions in order to use the current systems as far as possible in the event of an epidemic or lockdown
- Guidelines on provision of medical care to patients with NCDs through telemedicine technologies or alternate means such as ambulatory services or mHealth