



Highlights

Epidemiological week no. 51
(15 to 21 Dec 2013)

- Dengue fever:** During this week, 169 Dengue fever cases have been reported, out of them 145 from province Sindh; 20 from province Punjab; and 4 from province Balochistan. The outbreaks are being responded jointly.
- In this week, **77** districts and 2,278 health facilities have reported to Disease Early Warning System (DEWS), compared to 78 districts with 2,354 health facilities shared weekly data in week 50, 2013 to the DEWS.
- Total **789,224** patients consultations reported this week 51, 2013 compared with **896,479** consultations in week 50, 2013.
- 41** alerts investigated and **4** outbreaks identified and timely responded. Altogether 25 alerts were for Measles; 5 each for Dengue fever and Typhoid fever; 2 each for ARI and Leishmaniasis; while 1 each for Acute diarrhoea and Scabies.

Figure-1: 77 districts reported to DEWS in week 51, 2013



Priority diseases under surveillance in DEWS

Pneumonia
Acute Watery Diarrhoea
Bloody diarrhoea
Acute Diarrhoea
Suspected Enteric/Typhoid Fever
Suspected Malaria
Suspected Meningitis
Suspected Dengue fever
Suspected Viral Hemorrhagic Fever
Suspected Measles
Suspected Diphtheria
Suspected Pertussis
Suspected Acute Viral Hepatitis
Neonatal Tetanus
Acute Flaccid Paralysis
Scabies
Cutaneous Leishmaniasis


Cumulative number of selected health events reported in Epi-week 1 to 51, 2013 (29 Dec 2012 to 21 December 2013)

Disease	# of Cases	Percentage
ARI	8,182,886	20%
Bloody diarrhoea	98,035	<0.5%
Acute diarrhoea	3,004,031	7%
S. Malaria	1,743,762	4%
Skin Diseases	1,542,222	4%
Unexplained fever	1,251,718	3%
Total (All consultations)	40,916,821	

Major health events reported during the Epi-week - 51 (15 - 21 Dec 2013)


Disease	# of Cases	Percentage
ARI	172,390	22%
Bloody diarrhoea	1,303	<0.5%
Acute diarrhoea	41,506	5%
S. Malaria	21,794	3%
Skin Diseases	22,015	3%
Unexplained fever	22,560	3%
Total (All consultations)	789,224	

Figure-2: Number of Dengue fever positive cases in Pakistan, 1 Jan to 21 Dec 2013



From 1st January to 21st December 2013, Khyber Pakhtunkhwa reported 9,402; Sindh 5,645, Punjab 2,665, and Balochistan 42 confirmed Dengue fever cases.

In 2013 Dengue fever cases are reported from the less endemic areas also. A huge outbreak is confronted in district Swat and increasing number of Dengue fever cases are reported from adjacent district also and cases are also reported from Gawadar district in Balochistan province and Karachi in Sindh province.



Outbreaks (Wk-51/2013):

Date	Disease	Province	District	Area	<5M	>5M	<5F	>5F	Action Taken
16-Dec	Measles	Khyber Pakhtunkhwa	Battagram	Nowshera	0	0	5	0	Alert for 2 suspected Measles cases were reported from DHQ hospital Battagram. The cases belongs to district Nowshera. Vitamin-A drops given to all the suspected cases. In the past four weeks six suspected Measles cases have been reported from this area constituting a clinical outbreak. The EPI team requested to conduct outreach vaccination activity in the area. Information shared with DHO.
17-Dec	DF	Punjab	Lahore	Gulberg Town UC 99	0	3	0	5	1 confirmed, 6 suspected and 1 probable cases of Dengue fever reported from Teaching hospitals in Lahore. 1 blood sample taken and sent to laboratory testing and result found IgM positive. The confirmed case was admitted and kept in isolation ward (HDU) of Teaching hospital. There is no travel history of patient. Health education session given to the community on proper solid waste disposal and improved water storage practices, protective clothing and repellents, Community-based environmental management for improved water storage practices to remove mosquito breeding sites. Follow-up planned. Information shared with ADG(VBD) and EDO-H Lahore.
19-Dec	DF	Punjab	Lahore	Shalimar Town UC 24	0	6	0	1	2 confirmed and 5 suspected cases of Dengue fever notified from different teaching hospitals in Lahore. 2 blood samples were taken and sent to laboratory testing and result found IgM positive for both the cases. PCR request was sent for 1 case to IPH for serotyping and Genotyping, result is awaited. The confirmed cases were admitted and kept in isolation ward (HDU) of Teaching hospital. UC wise outbreak warning was generated by Dengue monitoring cell. Health education session was given to the community on proper solid waste disposal and improved water storage practices, protective clothing and repellents, Community-based environmental management for improved water storage practices to remove mosquito breeding sites. Follow-up planned. Information shared with ADG(VBD) and EDO-H Lahore.
21-Dec	DF	Punjab	Lahore	Shalimar Town UC 34	0	2	0	0	2 confirmed cases of Dengue fever notified from HDU Services Hospital Lahore. PCR request of patients were sent to PU for serotyping and genotyping result is awaited. The cases were admitted and kept in isolation ward (HDU) of Teaching hospital. There is no travel history found. DPC along with, CDDO, DDHO, and entomologist visit the area, Vector surveillance was carried out at confirmed case Location. Larva was found inside the houses of confirmed cases. Health education session given to the families and to the community on proper solid waste disposal and improved water storage practices, protective clothing and repellents and community-based environmental management on improved water storage practices to remove mosquito breeding sites. Follow-up planned. Information shared with ADG(VBD) and EDO-H Lahore.

Table-1: Number of alerts and outbreaks reported and investigated with appropriate response

Disease	2012		Current week 51, 2013		2013 (Total up till week - 51)	
	A	O	A	O	A	O
Acute watery diarrhoea	656	193	0	0	142	40
Acute jaundice syndrome	113	22	0	0	49	6
Bloody diarrhoea	146	11	0	0	44	3
CCHF	68	41	0	0	90	47
Dengue fever	175	29	5	3	300	66
Diphtheria	60	16	2	0	80	19
Measles	5922	812	25	1	3347	281
Pertussis	366	147	0	0	46	10
NNT + tetanus	560	0	0	0	344	0
Malaria	136	68	0	0	25	6
Cutaneous Leishmaniasis	900	78	0	0	613	49
Others	1529	58	9	0	509	5
Total	10631	1475	41	4	5589	532

This weekly Epidemiological Bulletin is published jointly by the National Institute of Health, Islamabad and World Health Organization (WHO), Pakistan.
For Correspondence: NIH: eic.nih@gmail.com; WHO: Tel : +92-051-9255184-5, Fax : +92-051-9255083; E-mail: wr@pak.emro.who.int.

Figure-3: Number of alerts received and responded, week 48 - 51, 2013






Figure-4: Number of consultations by age and gender, week 51, 2013



Province Khyber Pakhtunkhwa:

Figure-5: Weekly trend of Acute diarrhoea, province Khyber Pakhtunkhwa




201 health facilities from 9 districts of Khyber Pakhtunkhwa sent reports to DEWS with a total of 68,942 patients consultations reported in week 51, 2013. A total of 24 alerts for Measles were received and appropriate measures were taken.

The weekly trend of Acute diarrhoea is showing increase as compared with last week in KP, and 5 AWD confirmed cases reported from KP in last eight weeks, and the situation needs continuous attention in the province.

Province Sindh:

Figure-6: Weekly trend of Acute diarrhoea, province Sindh




806 health facilities from 23 districts in Sindh province reported to DEWS with a total of 228,690 patient consultations in week 51, 2013.

The proportion of AD for the province is showing minor increase as compared with last week, and during this season, 18 AWD outbreaks identified and responded, and the situation need continuous attention in the province.

Province Punjab:

Figure-7: Trend of Acute diarrhoea, province Punjab




888 health facilities from 18 districts in province Punjab reported to DEWS with a total of 423,647 patients consultations in week 51, 2013.

Total 12 alerts were received and appropriate measures were taken. Altogether 5 alerts were for Typhoid fever; 3 for Dengue fever; 2 for ARI; while 1 each for Acute diarrhoea and Scabies were responded in province Punjab.

The weekly trend of AD in Punjab showing minor decrease as compared with last two weeks, but required vigilant monitoring of the situation.

Province Balochistan:

Figure-8: Weekly trend of Acute diarrhoea, province Balochistan




269 health facilities from 14 districts in province Balochistan reported to DEWS with a total of 40,841 patients consultations in week 51, 2013.

5 alerts were reported and appropriated measures were taken. Altogether 2 alerts each were for Leishmaniasis and Dengue fever; while 1 for Measles.

In this week the weekly proportion of AD showing increase as compared with last week. and vigilant monitoring of the situation is required.

FATA:

Figure-9: Weekly trend of Acute diarrhoea, FATA




35 health facilities from 3 agencies in FATA reported to DEWS with a total of 12,929 patients consultations in week 51, 2013.

No alerts for any disease were reported from any area in FATA in week 51, 2013. But fluctuating and high weekly trend of Acute diarrhoea is noted in FATA and required vigilant monitoring.

State of Azad Jammu and Kashmir:

Figure-10: Weekly trend of Acute diarrhoea, AJ&K




72 health facilities from 8 districts in AJ&K reported to DEWS with a total of 12,069 patients consultations in week 51, 2013.

No alerts for any disease were reported from any area in AJ&K in week 51, 2013.

Weekly trend of Acute diarrhoea showing decrease as compared with last week, but vigilant monitoring of the situation is required.

Islamabad:

Figure-11: Weekly trend of Acute diarrhoea, Islamabad



6 health facilities reported to DEWS on time with a total of 1,925 patients consultations in week 51, 2013.

No alert for any disease was reported in week 51, 2013.


Weekly trend of Acute diarrhoea showing increase as compared with last week, and vigilant monitoring of the situation is required.

Distribution of Wild Polio Virus cases in Pakistan 2012 and 2013

In this week 51, 2013, two new type-1 wild polio cases were reported from Federally Administered Tribal Areas (North Waziristan Agency). This brings the total number of polio cases this year to 77 as compared to 58 last year, the number of infected districts/towns/tribal agencies/FR areas in the country is 20 as compared to 28 last year.

It is important to mention that 83% of total polio cases (64 cases) are from Khyber Pakhtunkhwa (10 cases) and FATA (54 cases). North Waziristan Agency in FATA, alone has reported 27 Wild Polio cases indicating very intense viral circulation in the tribal agency.

Province	2012			2013		
	P1	P3	P1+P3	P1	P3	P1+P3
Punjab	2	-	-	6	-	-
Sindh	4	-	-	7	-	-
Khyber Pakhtunkhwa	27	-	-	10	-	-
FATA	17	2	1	54	-	-
Balochistan	4	-	-	-	-	-
AJ&K	-	-	-	-	-	-
Gilgit-Baltistan	1	-	-	-	-	-
Islamabad	-	-	-	-	-	-
<i>Total</i>	<i>55</i>	<i>2</i>	<i>1</i>	<i>77</i>	<i>-</i>	<i>-</i>

Districts/ Towns with Wild Polio cases = 20**Follow up of CCHF**

In week 51, 2013, no new CCHF cases reported from any district. So for total of 100 suspected, 64 confirmed CCHF cases and 20 deaths have been reported country wide in year 2013.

In 2012, a total of 62 suspected cases were reported throughout the country with 41 cases confirmed and in total 18 deaths; of which 13 deaths (CFR is 31.7%) are reported of the lab confirmed cases and 5 deaths are reported as suspected CCHF cases. 23 confirmed cases were reported from Balochistan; 7 from Sindh; 6 from Khyber Pakhtunkhwa and 5 from Punjab. Table on right illustrates situation of CCHF cases in 2012-13.

Approximately all the cases had contact history with animal trading/handling, tick bite, contact with patient, tannery worker, butcher/animals slaughtering, and fresh animal skin. There is ongoing trade of animals and animal skins with movement intra Pakistan and between neighboring countries (Afghanistan and Iran).

Number of CCHF cases and deaths reported in year 2012 and 2013 up till 21 December.

Province	2012			2013		
	Suspected	Lab confirmed	Deaths	Suspected	Lab confirmed	Deaths
AJ&K	0	0	0	1	1	0
Balochistan	Afghanistan*	5	5	3	12	6
	Balochistan	33	18	4	54	33
ICT	-	-	-	4	4	1
KPK	9	6	5	9	9	4
Punjab	8	5	3	18	9	5
Sindh	7	7	3	2	2	1
Total	62	41	18	100	64	20

Focus on Dengue fever:

Dengue fever represents one of the most recent and emerging public health challenges with an ever-increasing incidence across the globe, way beyond its original impact area in the tropics. As estimated 2.5 billion people, are said to be at risk from this viral infection across the world. Dengue has emerged as a global problem after 1950's. Incidence of disease can be ascribed to a number of diverse factors related to hygiene, urban decay, environmental problems, changed living styles and physical factors. With a rapidly evolving epidemiology, the disease appears to have become more common in recent years and disturbing numbers of human mortalities have been reported.

The first confirmed outbreak of DF in Pakistan was in 1994 but the sudden rise in cases and annual epidemic trend occurred first in Karachi in November 2005. To assist with vector control, WHO entomologists studied the Aedes mosquito in Karachi during the 2005 outbreak and used the information to design Pakistan-specific control interventions. The worst ever epidemic of Dengue fever reported in Pakistan was in 2011, and reporting of cases started in August in Lahore – Punjab, with over 16,580 confirmed cases and 257 deaths reported from Lahore district and an additional 5000 cases and 60 deaths reported from the rest of the province.

In 2013 Dengue fever cases are reported from the less endemic areas also. A huge outbreak is confronted in district Swat and increasing number of Dengue fever cases are reported from adjacent district also and cases are also reported from Gawadar district in Balochistan province and Karachi in Sindh province. There is no vaccine available against dengue, and there are no specific medications to treat a dengue infection. This makes prevention the most important step, and prevention means avoiding mosquito bites if you live in or travel to an endemic area.

Patient management at home:

- If a person is having fever for more than 2 days with headache, retro-orbital pain, myalgia and arthralgia, it can be taken as a suspected dengue fever case and basic care should be started at home as follow
- Let the patient rest and reduce fever by sponging patient with tepid water. If the patient is shivering, stop sponging, and cover the patient with blanket. Give paracetamol every 4 hours only if the patient still has high fever $>39^{\circ}\text{C}$. **Do not give the patient Aspirin or Brufen or similar medicine.**
- If patient has any of the warning signs he/she should go to the hospital as soon as possible.

Warning signs (WS):

Following are the warning signs of the Dengue Hemorrhagic fever and require immediate referral to the hospital.

- Persistent vomiting, not drinking
- Severe abdominal pain, lethargy and restlessness
- Hematemesis, bleeding from gums and nose, excessive menstrual bleeding
- Hemoglobinuria, giddiness, pale, cold clammy hands and feet
- No/less urine for 4-6 hours


Prevention:

- The best way to reduce mosquitoes is to eliminate the places where the mosquito lays eggs, like water storage containers in and around the home, animal watering containers, flower planter dishes. Keep these containers empty or cover water storage barrels properly. Look for standing water indoors such as in vases with fresh flowers and clean at least once a week.
- The adult mosquitoes like to bite inside as well as around homes, during the day and at night when the lights are on. To protect yourself, wear long sleeves and pants, and use repellent on your skin while indoors or out.
- Use Mosquito coils, electric vapor mats during the day time. Curtains can also be treated with insecticides.
- Also, make sure window and door screens are secure and without holes. If available, use air-conditioning.
- If someone in your house is ill with dengue, take extra precautions to prevent mosquitoes from biting the patient and going on to bite others household members. Keep the dengue patient under the bed net.


Malaria:

The chart at below shows the Malaria slide positivity and Falciparum ratio in week 51, 2013. Total number of Malaria cases tested in this week is 7,191 out of which 740 were found positive; 405 for P. Vivax; 165 for P. Falciparum; while 170 for Mixed (SPR = 10.29%; FR = 45.27%).

Malaria tests \Province	Sindh	Punjab	Balochistan	KPK	FATA	AJK	GB	ICT
P. Vivax	96	28	104	115	62	0	0	0
P. Falcipa- rum	39	1	61	7	57	0	0	0
Mixed	139	0	6	2	23	0	0	0
# tested	2180	1909	1460	941	701	0	0	0
SPR	12.57	1.52	11.71	13.18	20.26	0	0	0
FR	64.96	3.45	39.18	7.26	56.34	0	0	0



Alerts and outbreaks, week 51, 2013



Produced by:
Health Information Management Unit,
WHO-Pakistan
Projection: WGS84
Data source: WHO (DEWS)

Website: www.emro.who.int/countries/pak

The boundaries and names shown and the designations used on this map do not imply the expression of any opinion whatsoever on the part of the World Health Organization concerning the legal status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries.
Dotted lines on maps represent approximate border lines for which there may not yet be full agreement.

GLIDE Number _____
WHO Map Num: WHO-PAK-1458
Date created: January 02, 2014

