

global youth tobacco survey

Country fact sheets

The World Health Organization (WHO) and the Centers for Disease Control and Prevention (CDC), Atlanta, developed the Global Youth Tobacco Survey to track tobacco use among youth across countries using a common methodology and core questionnaire. Information from the Survey is compiled within the participating country by a Research Coordinator nominated by the Ministry of Health, and technically reviewed by WHO and CDC. The content has not otherwise been edited by WHO or CDC.

All reasonable precautions have been taken by the World Health Organization to verify the information contained in this publication. However, the published material is being distributed without warranty of any kind, either expressed or implied. The responsibility for the interpretation and use of the material lies with the reader. In no event shall the World Health Organization be liable for damages arising from its use.



**World Health
Organization**

Regional Office for the Eastern Mediterranean



Jordan 2009 (Ages 13-15) Global Youth Tobacco Survey (GYTS) FACT SHEET



The Jordan GYTS includes data on prevalence of cigarette and other tobacco use as well as information on five determinants of tobacco use: access/availability and price, exposure to secondhand smoke (SHS), cessation, media and advertising, and school curriculum. These determinants are components Jordan could include in a comprehensive tobacco control program.

The Jordan GYTS was a school-based survey of students in grades 7th, 8th, and 9th conducted in 2009.

A two-stage cluster sample design was used to produce representative data for Jordan. At the first stage, schools were selected with probability proportional to enrollment size. At the second stage, classes were randomly selected and all students in selected classes were eligible to participate. A total of 2,064 students participated in the Jordan GYTS of which 1,523 were ages 13 to 15 years. The overall response rate of all students surveyed was 90.3%.

Prevalence

- 25.2% of students had ever smoked cigarettes (Boys = 34.8%, Girls = 17.8%)
- 26.1% currently use any tobacco product (Boys = 34.1%, Girls = 19.4%)
- 11.5% currently smoke cigarettes (Boys = 17.4%, Girls = 6.6%)
- 20.7% currently smoke shisha (Boys = 27.1%, Girls = 15.6%)
- 22.1% currently use other tobacco products (Boys = 28.4%, Girls = 16.7%)
- 23.7% of never smokers are likely to initiate smoking next year

Knowledge and Attitudes

23.7% think boys and 16.7% think girls who smoke look more attractive

Access and Availability - Current Smokers

- 43.4% buy cigarettes in a store
- 93.4% who bought cigarettes in a store were NOT refused purchase because of their age

Exposure to Secondhand Smoke (SHS)

- 53.6% live in homes where others smoke in their presence
- 50.5% are around others who smoke in places outside their home
- 76.5% think smoking should be banned from public places
- 59.4% think smoke from others is harmful to them
- 49.4% have one or more parents who smoke
- 16.4% have most or all friends who smoke

Cessation - Current Smokers

- 51.6% want to stop smoking
- 63.6% tried to stop smoking during the past year
- 73.6% have ever received help to stop smoking

Media and Advertising

- 69.1% saw anti-smoking media messages, in the past 30 days
- 61.9% saw pro-cigarette ads in newspapers or magazines, in the past 30 days
- 18.0% have an object with a cigarette brand logo
- 11.0% were offered free cigarettes by a tobacco company representative

School

- 39.0% had been taught in class, during the past year, about the dangers of smoking
- 43.6% had discussed in class, during the past year, reasons why people their age smoke

Highlights

- 26.1% of students currently use any form of tobacco; 11.5% currently smoke cigarettes; 20.7% currently smoke shisha; 22.1% currently use some other form of tobacco
- SHS exposure – 53.6% of students live in homes where others smoke, and half the students are exposed to smoke around others outside of the home; half the students have at least one parent who smokes
- Three in 5 students think smoke from others is harmful to them
- Over 4 in 5 students think smoking in public places should be banned
- More than half the current smokers want to stop smoking
- 18.0% of students have an object with a cigarette brand logo on it
- Seven in 10 students saw anti-smoking media messages in the past 30 days; over 6 in 10 saw pro-cigarette ads in newspapers or magazines in the past 30 days