

Country fact sheets

The World Health Organization (WHO) and the Centers for Disease Control and Prevention (CDC), Atlanta, developed the Global Youth Tobacco Survey to track tobacco use among youth across countries using a common methodology and core questionnaire. Information from the Survey is compiled within the participating country by a Research Coordinator nominated by the Ministry of Health, and technically reviewed by WHO and CDC. The content has not otherwise been edited by WHO or CDC.

All reasonable precautions have been taken by the World Health Organization to verify the information contained in this publication. However, the published material is being distributed without warranty of any kind, either expressed or implied. The responsibility for the interpretation and use of the material lies with the reader. In no event shall the World Health Organization be liable for damages arising from its use.



Regional Office for the Eastern Mediterranean



© World Health Organization 2015

GYTS Objectives

The Global Youth Tobacco Survey (GYTS), a component of the Global Tobacco Surveillance System (GTSS), is a global standard for systematically monitoring youth tobacco use (smoking and smokeless) and tracking key tobacco control indicators.

GYTS is a nationally representative school-based survey of students in grades associated with 13 to 15 years of age and is designed to produce cross-sectional estimates for each country. GYTS uses a standard core questionnaire, sample design, and data collection protocol. It assists countries in fulfilling their obligations under the World Health Organization (WHO) Framework Convention on Tobacco Control (FCTC) to generate comparable data within and across countries. WHO has developed MPOWER, a package of selected demand reduction measures contained in the WHO FCTC:

mpow

Monitor tobacco use & prevention policies

Protect people from tobacco smoke

Offer help to guit tobacco use

Warn about the dangers of tobacco

Enforce bans on tobacco advertising, promotion, & sponsorship

Raise taxes on tobacco

GYTS Methodology

GYTS uses a global standardized methodology that includes a two-stage sample design with schools selected with a probability proportional to enrollment size. The classes within selected schools are chosen randomly and all students in selected classes are eligible to participate in the survey. The survey uses a standard core questionnaire with a set of optional questions that permits adaptation to meet the needs of the country on tobacco use and key tobacco control indicators. The questionnaire consists of the following topics: tobacco use (smoking and smokeless), cessation, secondhand smoke (SHS), pro- and antitobacco media and advertising, access and availability to obtain tobacco products, and knowledge and attitudes regarding tobacco. The questionnaire is self-administered; using scannable paper-based bubble sheets, it is anonymous to ensure confidentiality.

In UNRWA West Bank, GYTS was conducted in 2014 by UNRWA Headquarters, under the coordination of the Health and Education Departments. A total of 1,367 eligible students in grades 7-9 completed the survey, of which 1,172 were aged 13-15 years. The overall response rate of all students surveyed was 93.4%.

GYTS Highlights

TOBACCO USE

- 34.6% overall, 47.5% of boys, and 24.7% of girls currently • used any tobacco products.
- 31.8% overall, 44.8% of boys, and 22.0% of girls currently smoked tobacco.
- 21.0% overall, 31.2% of boys, and 13.1% of girls currently smoked cigarettes.
- 4.8% overall, 5.8% of boys, and 4.1% of girls currently used smokeless tobacco.
- 19.2% overall, 30.8% of boys, and 9.7% of girls currently smoked electronic cigarettes.

CESSATION

- More than 5 in 10 current smokers tried to stop smoking in the past 12 months.
- Almost 6 in 10 current smokers want to stop smoking now.

SECONDHAND SMOKE

- 69.0% of students were exposed to tobacco smoke at home.
- 66.7% of students were exposed to tobacco smoke inside enclosed public places.

ACCESS & AVAILABILITY

- 42.2% of current cigarette smokers obtained cigarettes by buying them from a store, shop, street vendor, or kiosk.
- Among current cigarette smokers who bought cigarettes, 72.4% were not prevented from buying them because of their age.

MEDIA

- 6 in 10 students noticed anti-tobacco messages in the media.
- 5 in 10 students noticed tobacco advertisements or promotions when visiting points of sale.
- Almost 2 in 10 students own something with a tobacco brand logo on it.

KNOWLEDGE & ATTITUDES

- 72.1% of students definitely thought other people's tobacco smoking is harmful to them.
- 73.8% of students favor banning smoking inside enclosed public places.










GYTS GLOBAL YOUTH TOBACCO SURVEY

FACT SHEET UNRWA WEST BANK 2014

TOBACCO USE

SMOKED TOBACCO	OVERALL (%)	BOYS (%)	GIRLS (%)
Current tobacco smokers ¹	31.8	44.8	22.0
Current cigarette smokers ²	21.0	31.2	13.1
Frequent cigarette smokers ³	3.8	8.8	0.0
Current smokers of other tobacco ⁴	21.8	30.5	15.3
Ever tobacco smokers ⁵	61.2	72.6	52.4
Ever cigarette smokers ⁶	51.9	62.7	43.8
Ever smokers of other tobacco ⁷	43.4	53.0	36.2
SMOKELESS TOBACCO			
Current smokeless tobacco users ⁸	4.8	5.8	4.1
Ever smokeless tobacco users ⁹	9.2	10.7	7.9
TOBACCO USE (smoked and/or smokeless)			
Current tobacco users ¹⁰	34.6	47.5	24.7
Ever tobacco users ¹¹	63.9	74.2	55.8
SUSCEPTIBILITY			
Never to bacco users susceptible to to bacco use in the $\rm future^{12}$	10.8	17.7	7.8
Never smokers who thought they might enjoy smoking a cigarette ¹³	8.5	11.8	6.9



CESSATION

	OVERALL (%)	BOYS (%)	GIRLS (%)
Current smokers who tried to stop smoking in the past 12 months	54.1	58.4	45.0
Current smokers who want to stop smoking now	56.7	58.9	51.0
Current smokers who thought they would be able to stop smoking if they wanted to	68.7	63.8	79.1
Current smokers who have ever received help/advice from a program or professional to stop smoking	26.9	24.1	31.8

SECONDHAND SMOKE

	OVERALL (%)	BOYS (%)	GIRLS (%)
Exposure to tobacco smoke at home ⁺⁺	69.0	66.8	70.9
Exposure to tobacco smoke inside any enclosed public place ^{††}	66.7	72.5	62.0
Exposure to tobacco smoke at any outdoor public place ^{††}	64.8	69.2	61.4
Students who saw anyone smoking inside the school building or outside on school property [†]	54.9	67.8	45.2

ACCESS & AVAILABILITY

	OVERALL (%)	BOYS (%)	GIRLS (%)
Current cigarette smokers who obtained cigarettes by buying them from a store, shop, street vendor, or kiosk ¹⁴	42.2	61.0	8.5
Current cigarette smokers who were not prevented from buying cigarettes because of their age ¹⁵	72.4	70.7	
Current cigarette smokers who bought cigarettes as individual sticks ¹⁶	47.5	47.4	
MEDIA			
TOBACCO INDUSTRY ADVERTISING	OVERALL (%)	BOYS (%)	GIRLS (%)
Noticing tobacco advertisements or promotions at points of sale ¹⁷	52.0	60.2	45.7
Students who saw anyone using tobacco on television, videos, or movies ¹⁸	85.0	82.0	87.0
Students who were ever offered a free tobacco product from a tobacco company representative	12.5	18.7	7.9
Students who own something with a tobacco brand logo on it	16.8	23.0	12.3
ANTI-TOBACCO ADVERTISING			
Noticing anti-tobacco messages in the media [†]	63.4	60.6	65.4
Noticing anti-tobacco messages at sporting or community events ¹⁹	49.7	49.8	49.0
Current smokers who thought about quitting because of a warning label ²⁰	36.4	39.7	31.3
Students who were taught in school about the dangers of tobacco use in the past 12 months	62.0	55.6	67.2

KNOWLEDGE & ATTITUDES

	OVERALL (%)	BOYS (%)	GIRLS (%)
Students who definitely thought it is difficult to quit once someone starts smoking tobacco	34.9	31.4	37.7
Students who thought smoking tobacco helps people feel more comfortable at celebrations, parties, and social gatherings	22.4	27.1	18.9
Students who definitely thought other people's tobacco smoking is harmful to them	72.1	64.7	78.2
Students who favor banning smoking inside enclosed public places	73.8	65.9	79.8
Students who favor banning smoking at outdoor public places	66.1	61.2	70.0

¹ Smoked tobacco anytime during the past 30 days. ² Smoked cigarettes anytime during the past 30 days. ⁵ Smoked cigarettes on 20 or more days of the past 30 days. ⁴ Smoked tobacco other than cigarettes anytime during the past 30 days. ⁵ Ever smoked any tobacco, even one or two puffs. ⁶ Ever smoked tobacco other than cigarettes, even one or two puffs. ⁸ Used smokeless tobacco anytime during the past 30 days. ⁹ Ever used smokeless tobacco. ¹⁰ Smoked tobacco other than cigarettes, even one or two puffs. ⁸ Used smokeless tobacco anytime during the past 30 days. ⁹ Ever used smokeless tobacco. ¹² Smoked tobacco and/or used smokeless tobacco. ¹² Susceptible to future tobacco is encludes those who answered "Definitely yes", "Probably yes", or "Probably not" to using tobacco if one of their best friends offered it to them or those who answered "Definitely yes", "Probably yes", or "Probably not" to using tobacco if one of their best friends offered it to them or those who answered "Definitely yes", "Probably yes", or "Probably not" to using tobacco if one of their best friends offered it to them or those who answered "Definitely yes", "Probably yes", or "Probably not" to using tobacco if one of their best friends offered it to them or those who answered "Definitely yes", "Probably yes", or "Brobably not" to using tobacco and/or used smokeless tobacco. ¹⁵ Of those who cigarettes user obtained the last time respondents smoked cigarettes in the past 30 days. ¹⁵ Of those who tried to buy cigarettes during the past 30 days. ¹⁶ Based on the last purchase, of those who bught cigarettes during the past 30 days. ¹⁹ Among those who noticed warning labels on cigarette packages in the past 30 days. ²⁰ Among those who noticed warning labels on cigarette packages in the past 30 days. ¹⁵ During the past 30 days. ¹⁵ During the past 30 days. ¹⁵ During the past 30 days.

NOTE: Students refer to persons aged 13-15 years who are enrolled in school. Data have been weighted to be nationally representative of all students aged 13-15 years. Percentages reflect the prevalence of each indicator in each group, not the distribution across groups.

--Indicates estimate based on less than 35 unweighted cases and has been suppressed.