

# global health professions student survey

## Country fact sheets

The World Health Organization (WHO) and the Centers for Disease Control and Prevention (CDC), Atlanta, developed the Global Health Professions Student Survey to track tobacco use among health professions students across countries using a common methodology and core questionnaire. Information from the Survey is compiled within the participating country by a Research Coordinator nominated by the Ministry of Health, and technically reviewed by WHO and CDC. The content has not otherwise been edited by WHO or CDC.

All reasonable precautions have been taken by the World Health Organization to verify the information contained in this publication. However, the published material is being distributed without warranty of any kind, either expressed or implied. The responsibility for the interpretation and use of the material lies with the reader. In no event shall the World Health Organization be liable for damages arising from its use.



**World Health  
Organization**

Regional Office for the Eastern Mediterranean



**CENTERS FOR DISEASE  
CONTROL AND PREVENTION**

# Saudi Arabia - Dental Students 2010 (3<sup>rd</sup> Year Students Only) Global Health Professions Student Survey (GHPSS)

## FACT SHEET . . . . .

The Saudi Arabia Dental students GHPSS includes data on prevalence of cigarette smoking and use of other tobacco products, knowledge and attitudes about tobacco use, exposure to second-hand smoke, desire for smoking cessation, and training received during Dental school regarding patient counseling on smoking cessation techniques. This information could be very useful in monitoring and developing Saudi Arabia's comprehensive tobacco control program.

The Saudi Arabia Dental students GHPSS was a census of 3<sup>rd</sup> year Dental students attending 5 Dental schools in Saudi Arabia conducted in 2010. A total of 73 3<sup>rd</sup> year students participated in the Saudi Arabia Dental students GHPSS. The overall response rate of all Dental students surveyed was 83.1%.

### Prevalence

- 47.4% ever smoked cigarettes (Males = 64.8%, Females = 25.3%)
- 25.9% currently smoke cigarettes (Males = 40.1%, Females = 9.0%)
- 54.6% ever used any form of tobacco other than cigarettes (Males = 67.9%, Females = 38.9%)
- 34.3% currently use any form of tobacco other than cigarettes (Males = 54.1%, Females = 9.4%)
- 34.3% currently smoke shisha (Male = 54.1%, Female = 9.4%)
- 31.3% of ever smokers smoked on school premises/property during the past year<sup>†</sup>
- 10.9% of ever smokers smoked in school buildings during the past year<sup>†</sup>

### Factors Influencing Tobacco Use

- 37.2% were exposed to second-hand smoke at home, during the past week
- 61.4% were exposed to second-hand smoke in public places during the past week
- 68.2% reported that their school had a ban on smoking in school buildings and clinics
- 95.2% reported that their school enforced the ban on smoking in school buildings and clinics
- 92.1% thought smoking should be banned in all enclosed public places
- 92.9% thought tobacco sales to adolescents should be banned
- 92.0% thought there should be a complete ban on advertising tobacco products.
- 80.0% of current smokers wanted to stop smoking<sup>†</sup>
- 50.7% of current smokers ever received help/advice to stop smoking cigarettes<sup>†</sup>
- 85.8% were taught about the dangers of smoking during dental school training

### Role Model and Cessation Training

- 83.3% thought health professionals serve as role models for their patients
- 89.6% thought health professionals have a role in giving advice about smoking cessation to patients
- 19.3% received formal training in smoking cessation approaches during dental school
- 90.5% thought health professionals should get specific training on cessation techniques

(<sup>†</sup> The percentage is based on observations between 10 and 34)

### Highlights

**25.9% currently smoke cigarettes and 34.9% currently use other tobacco products**

**Eight in 10 smokers indicated that they want to quit**

**Six in 10 were exposed to smoke in public places**

**Over nine in 10 support a ban on smoking in all enclosed public places**

**Two in 10 received training in smoking cessation approaches to use with patients**

**Nine in 10 think health professionals should get specific training on cessation techniques to use with patients**