## **GYTS Objectives**

The Global Youth Tobacco Survey (GYTS), a component of the Global Tobacco Surveillance System (GTSS), is a global standard for systematically monitoring youth tobacco use (smoking and smokeless) and tracking key tobacco control indicators.

GYTS is a cross-sectional, nationally representative school-based survey of students in grades associated with ages 13 to 15 years. GYTS uses a standard core questionnaire, sample design, and data collection protocol. It assists countries in fulfilling their obligations under the World Health Organization (WHO) Framework Convention on Tobacco Control (FCTC) to generate comparable data within and across countries. WHO has developed MPOWER, a technical package of selected demand reduction measures contained in the WHO FCTC:



Monitor tobacco use & prevention policies

Protect people from tobacco smoke

Offer help to quit tobacco use

Warn about the dangers of tobacco

Enforce bans on tobacco advertising, promotion, & sponsorship

Raise taxes on tobacco

# **GYTS Methodology**

GYTS uses a global standardized methodology that includes a two-stage sample design with schools selected with a probability proportional to enrollment size. The classes within selected schools are chosen randomly and all students in selected classes are eligible to participate in the survey. The survey uses a standard core questionnaire with a set of optional questions that countries can adapt to measure and track key tobacco control indicators. The questionnaire covers the following topics: tobacco use (smoking and smokeless), cessation, secondhand smoke (SHS), pro- and anti-tobacco media and advertising, access to and availability of tobacco products, and knowledge and attitudes regarding tobacco use. The questionnaire is self-administered; using scannable paper-based bubble sheets, it is anonymous to ensure confidentiality.

In the Islamic Republic of Iran, GYTS was conducted in 2016 by the Ministry of Health and Medical Education. The overall response rate was 94.4%. A total of 2,746 eligible students in 2nd and 3rd grades of secondary school and first grade of high school (grades 8-10) completed the survey, of which 2,087 were aged 13-15 years. Data are reported for students aged 13-15 years.



### **GYTS Highlights**

#### **TOBACCO USE**

- 10.2% of students, 12.9% of boys, and 7.7% of girls currently used any tobacco products.
- 9.2% of students, 11.6% of boys, and 7.1% of girls currently smoked tobacco.
- 3.4% of students, 4.8% of boys, and 2.1% of girls currently smoked cigarettes.
- 11.2% of students, 14.3% of boys, and 8.5% of girls currently smoked shisha.
- 1.9% of students, 3.1% of boys, and 0.8% of girls currently used smokeless tobacco.

#### **CESSATION**

- More than 4 in 10 (41.4%) current tobacco smokers tried to stop smoking in the past 12 months.
- More than 5 in 10 (53.5%) current tobacco smokers wanted to stop smoking now.

### **SECONDHAND SMOKE**

- 24.2% of students were exposed to tobacco smoke at home.
- 38.4% of students were exposed to tobacco smoke inside enclosed public places.

#### **ACCESS & AVAILABILITY**

- 60.5% of current cigarette smokers bought cigarettes from a shop, supermarket, street vendor, or newspaper booth.
- Among current cigarette smokers who tried to buy cigarettes, 68.5% were not prevented from buying them because of their age.

#### **MEDIA**

- Almost 7 in 10 (69.9%) students noticed anti-tobacco messages in the media.
- More than 4 in 10 (40.2%) students noticed tobacco advertisements or promotions when visiting points of sale.
- More than 1 in 10 (11.8%) students had something with a tobacco brand logo on it.


#### **KNOWLEDGE & ATTITUDES**

- 76.7% of students definitely thought other people's tobacco smoking is harmful to them.
- 78.2% of students favored prohibiting smoking inside enclosed public places.



# **ISLAMIC REPUBLIC OF IRAN 2016**

TOBACCO USE			
SMOKED TOBACCO	OVERALL (%)	BOYS (%)	GIRLS (%)
Current tobacco smokers <sup>1</sup>	9.2	11.6	7.1*
Current cigarette smokers <sup>2</sup>	3.4	4.8	2.1*
Frequent cigarette smokers <sup>3</sup>	0.8	1.4	0.2*
Current smokers of other tobacco <sup>4</sup>	7.8	9.7	6.1*
Current shisha smokers <sup>5</sup>	11.2	14.3	8.5*
Frequent shisha smokers <sup>6</sup>	1.2	1.7	0.8
Ever tobacco smokers <sup>7</sup>	29.5	35.4	24.1*
Ever cigarette smokers <sup>8</sup>	14.2	16.8	11.8*
Ever smokers of other tobacco <sup>9</sup>	24.2	28.5	20.2*
Ever shisha smokers <sup>10</sup>	35.8	42.1	30.2*
SMOKELESS TOBACCO			
Current smokeless tobacco users <sup>11</sup>	1.9	3.1	0.8*
Ever smokeless tobacco users <sup>12</sup>	2.6	4.1	1.2*
ANY TOBACCO USE (smoked and/or smokeless)			
Current tobacco users <sup>13</sup>	10.2	12.9	7.7*
Ever tobacco users <sup>14</sup>	30.4	36.7	24.6*
SUSCEPTIBILITY			
Never to bacco users susceptible to to bacco use in the future $^{\rm 15}$	10.1	7.5	12.1*
Never to bacco smokers who thought they might enjoy smoking a cigarett e $^{\rm 16}$	6.4	5.2	7.3



### **CESSATION**

	OVERALL (%)	BOYS (%)	GIRLS (%)
Current tobacco smokers who tried to stop smoking in the past 12 months	41.4	44.4	36.9
Current tobacco smokers who wanted to stop smoking now	53.5	62.7	
Current tobacco smokers who thought they would be able to stop smoking if they wanted to	74.3	73.4	75.7
Current tobacco smokers who have ever received help/advice from a program or professional to stop smoking	26.8	25.1	29.2

#### **SECONDHAND SMOKE**

	OVERALL (%)	BOYS (%)	GIRLS (%)
Exposure to tobacco smoke at home <sup>17</sup>	24.2	26.9	21.7
Exposure to tobacco smoke inside any enclosed public place <sup>17</sup>	38.4	40.3	36.7
Exposure to tobacco smoke at any outdoor public place <sup>17</sup>	42.2	43.4	41.0
Students who saw anyone smoking inside the school building or outside on school property $^{18}$	28.7	36.3	21.6*

I	ACCESS & AVAILABILITY				
		OVERALL (%)	BOYS (%)	GIRLS (%)	
	Current cigarette smokers who bought cigarettes from a shop, supermarket, street vendor, or newspaper booth $^{19}$	60.5			
	Current cigarette smokers who were not prevented from buying cigarettes because of their age <sup>20</sup>	68.5			
	Current cigarette smokers who bought cigarettes as individual sticks <sup>21</sup>	41.8			

MEDIA			
TOBACCO ADVERTISING	OVERALL (%)	BOYS (%)	GIRLS (%)
Students who noticed to bacco advertisements or promotions at points of sale $^{\rm 22}$	40.2	41.7	38.7
Students who saw anyone using tobacco on television, videos, or movies <sup>23</sup>	61.8	60.4	63.1
Students who were ever offered a free tobacco product from a tobacco company representative	7.7	11.7	4.2*
Students who had something with a tobacco brand logo on it	11.8	13.8	10.0
ANTI-TOBACCO ADVERTISING			
Students who noticed anti-tobacco messages in the media $^{18}$	69.9	70.2	69.6
Students who noticed anti-tobacco messages at sporting or community events $^{24}$	36.8	43.9	28.6*
Current tobacco smokers who thought about quitting because of a warning label <sup>25</sup>	22.0	25.2	17.5
Students who were taught in school about the dangers of tobacco use in the past 12 months	38.0	42.7	33.8

KNOWLEDGE & ATTITUDES			
	OVERALL (%)	BOYS (%)	GIRLS (%)
Students who definitely thought it is difficult to quit once someone starts smoking tobacco	37.3	34.3	40.0*
Students who thought smoking tobacco helps people feel more comfortable at ceremonies or other social gatherings	24.1	28.3	20.5*
Students who definitely thought other people's tobacco smoking is harmful to them	76.7	72.8	80.2
Students who favored prohibiting smoking inside enclosed public places	78.2	73.9	82.0
Students who favored prohibiting smoking at outdoor public places	74.8	70.4	78.8*

<sup>1</sup> Smoked cigarettes or other type of tobacco anytime during the past 30 days. <sup>2</sup> Smoked cigarettes anytime during the past 30 days. <sup>3</sup> Smoked cigarettes on 20 or more days of the past 30 days. <sup>4</sup> Smoked tobacco other than cigarettes anytime during the past 30 days. <sup>5</sup> Smoked shisha anytime during the past 30 days. <sup>6</sup> Smoked shisha on 20 or more days of the past 30 days. <sup>7</sup> Ever smoked cigarettes or other type of tobacco, even one or two puffs. <sup>10</sup> Ever smoked cigarettes, even one or two puffs. <sup>10</sup> Ever smoked cigarettes, even one or two puffs. <sup>11</sup> Used smokeless tobacco anytime during the past 30 days. <sup>12</sup> Ever used smokeless tobacco. <sup>13</sup> Smoked cigarettes, smoked other type of tobacco, and/or used smokeless tobacco anytime during the past 30 days. <sup>14</sup> Ever smoked tobacco and/or used smokeless tobacco. <sup>15</sup> Susceptible to future tobacco use includes those who answered "definitely yes", "probably yes", or "probably not" to using tobacco if one of their best friends offered it to them, or "definitely yes", "probably yes", or "probably not" to using tobacco during the next 12 months. <sup>16</sup> Those who answered "Agree" or "Strongly Agree" to the statement: "I think I might enjoy smoking a cigarette". <sup>17</sup> During the past 7 days. <sup>18</sup> During the past 30 days. <sup>20</sup> Among those who tried to buy cigarettes the last time they smoked cigarettes in the past 30 days. <sup>22</sup> Among those who bought cigarettes during the past 30 days. <sup>22</sup> Based on the last purchase, among those who bought cigarettes during the past 30 days. <sup>23</sup> Among those who watched television, videos, or movies in the past 30 days. <sup>24</sup> Among those who attended sporting or community events in the past 30 days. <sup>25</sup> Among those who noticed warning labels on cigarette packages in the past 30 days. <sup>26</sup> Among those of community events in the past 30 days. <sup>28</sup> Among those who noticed warning labels on cigarette packages in the past 30 days.

**NOTE:** Data are weighted to be nationally representative of all students aged 13-15 years who are enrolled in school. Percentages reflect the prevalence of each indicator in each group, not the distribution across groups. Shisha estimates are not included in the tobacco use indicators.

<sup>--</sup>Estimates based on unweighted cases less than 35 are not presented.

<sup>\*</sup>Gender comparisons are significant at p < 0.05.