

Country fact sheets

The World Health Organization (WHO) and the Centers for Disease Control and Prevention (CDC), Atlanta, developed the Global Youth Tobacco Survey to track tobacco use among youth across countries using a common methodology and core questionnaire. Information from the Survey is compiled within the participating country by a Research Coordinator nominated by the Ministry of Health, and technically reviewed by WHO and CDC. The content has not otherwise been edited by WHO or CDC.

All reasonable precautions have been taken by the World Health Organization to verify the information contained in this publication. However, the published material is being distributed without warranty of any kind, either expressed or implied. The responsibility for the interpretation and use of the material lies with the reader. In no event shall the World Health Organization be liable for damages arising from its use.



Regional Office for the Eastern Mediterranean



© World Health Organization 2010

media and advertising, and school curriculum. These determinants are components Libya could include in a comprehensive tobacco control program. The Libya GYTS was a school-based survey of students in grades 7, 8 and 9 conducted in 2003. A two-stage cluster were eligible to participate. The the student response rate was 9 rate was 97%. A total of 1,174 participated in the Libya GYTS

sample design was used to produce representative data for all of Libya. At the first stage, schools were selected with probability proportional to enrollment size. At the second stage, classes were randomly selected and all students in selected classes were eligible to participate. The school response rate was 98%, the student response rate was 99%, and the overall response rate was 97%. A total of 1,174 students aged 13-15 participated in the Libya GYTS.

Prevalence

12.3% of students had ever smoked cigarettes (Boy = 19.2%, Girl = 5.2%)

12.7% currently use any tobacco product (Boy = 16.8%, Girl = 8.1%) 4.1% currently smoke cigarettes (Boy = 7.3%, Girl = 0.8%)

The Libya GYTS includes data on prevalence of cigarette

determinants of tobacco use: access/availability and price,

environmental tobacco smoke exposure (ETS), cessation,

and other tobacco use as well as information on five

- 4.1% currently smoke cigarettes (Boy = 7.5%, Girl = 0.8%) 9.8% currently use other tobacco products (Boy = 11.7%, Girl = 7.5%)
- 9.8% currently use other tobacco products (Boy = 11.7%, Girl = 7.5%19.5% of never smokers are likely to initiate smoking next year

Knowledge and Attitudes

26.0% think boys and 18.6% think girls who smoke have more friends 17.5% think boys and 13.1% think girls who smoke look more attractive

Access and Availability - Current Smokers

22.5% usually smoke at home 19.4% buy cigarettes in a store

Environmental Tobacco Smoke

40.4% live in homes where others smoke in their presence 38.6% are around others who smoke in places outside their home 77.3% think smoking should be banned from public places 69.0% think smoke from others is harmful to them 31.4% have one or more parents who smoke

4.7% have most or all friends who smoke

Cessation - Current Smokers

88.8% have ever received help to stop smoking

Media and Advertising

- 70.0% saw anti-smoking media messages, in the past 30 days
- 54.6% saw pro-cigarette ads on billboards, in the past 30 days
- 50.6% saw pro-cigarette ads in newspapers or magazines, in the past 30 days
- 10.1% have an object with a cigarette brand logo
- 8.2% were offered free cigarettes by a tobacco company representative

School

51.5% had been taught in class, during the past year, about the dangers of smoking 34.7% had discussed in class, during the past year, reasons why people their age smoke 43.7% had been taught in class, during the past year, the effects of tobacco use

Highlights

- 12.7% of students currently use any form of tobacco; 4.1% currently smoke cigarettes; 9.8% currently use some other form of tobacco.
- ETS exposure is high Over 4 in 10 students live in homes where others smoke in their presence; Almost 4 in 10 are exposed to smoke in public places; Over half have one or more parents who smoke.
- Over two-thirds of the students think smoke from others is harmful to them.
- More than three-quarters of the students think smoking in public places should be banned.
- Seven in 10 students saw antismoking media messages in the past 30 days; More than half the students saw pro-cigarette ads on billboards in the past 30 days; Over 5 in 10 saw pro-cigarette ads in newspapers or magazines in the past 30 days.
- One in 10 students has an object with a cigarette brand logo on it.



