

[High heterogeneity of malaria transmission and a large sub-patent and diverse reservoir of infection in Wusab As Safil district, Republic of Yemen](#)

[Susceptibility tests on insecticides used to control mosquitoes in Jordan](#)

[Effectiveness and cost of insecticide-treated bed nets and indoor residual spraying for the control of cutaneous leishmaniasis: a cluster-randomized control trial in Morocco](#)

Evidence of metabolic mechanisms playing a role in multiple insecticides resistance in *Anopheles stephensi* populations from Afghanistan

Design of a study to determine the impact of insecticide resistance on malaria vector control: a multi-country investigation

[Distribution and bionomic of sand flies in five ecologically different cutaneous leishmaniasis foci in Morocco](#)

[Insecticide susceptibility status of *Phlebotomus* \(*Paraphlebotomus*\) *sergenti* and *Phlebotomus* \(*Phlebotomus*\) *papatasi* in endemic foci of cutaneous leishmaniasis in Morocco](#)

[Implementation of integrated vector management for disease vector control in the Eastern Mediterranean where are we and where are we going? Eastern Mediterranean Health Journal Vol. 17 No. 5 2011](#)

[Management of the use of public health pesticides in the face of the increasing burden of vector-borne diseases in the Eastern Mediterranean region. Eastern Mediterranean Health Journal Vol. 18 No. 1 2012](#)

Saturday 17th of May 2025 02:12:41 PM