

Table 2 Changes in self-reported DASS-21 pre-and post-intervention scores in the music intervention and control (no intervention) groups, before and after angiography

| DASS-21 item | Before angiography | | | | | After angiography | | | | |
|-------------------|--------------------|--------------|--------------------|----------------------|---------|---------------------|-------------|--------------------|-----------------------|---------|
| | Control group | | Intervention group | Pre-/post-comparison | | Control group | | Intervention group | Pre- /post-comparison | |
| | Mean (SD) | Mean (SD) | | t-test ^a | P-value | Difference in means | Mean (SD) | Mean (SD) | t-test ^a | P-value |
| Stress score | | | | -3.51 | 0.001 | -2.89 | | | -1.49 | 0.14 |
| Pre-intervention | 11.00 (4.94) | 11.35 (4.31) | | | | | 5.89 (5.15) | 3.91 (4.44) | | |
| Post-intervention | 8.70 (4.89) | 6.16 (4.98) | | | | | 5.54 (4.43) | 4.37 (4.32) | | |
| Anxiety score | | | | -2.83 | 0.006 | -2.00 | | | -0.06 | 0.94 |
| Pre-intervention | 6.81 (4.29) | 6.83 (4.14) | | | | | 4.32 (4.23) | 2.83 (3.55) | | |
| Post-intervention | 6.10 (3.71) | 4.13 (3.81) | | | | | 3.81 (3.23) | 2.35 (2.46) | | |
| Depression score | | | | -2.37 | 0.02 | -1.35 | | | -0.69 | 0.48 |
| Pre-intervention | 5.86 (4.19) | 6.59 (4.55) | | | | | 4.72 (5.00) | 4.05 (4.75) | | |
| Post-intervention | 5.02 (3.79) | 4.40 (4.42) | | | | | 4.67 (3.59) | 3.70 (3.55) | | |

^adf = 72.

DASS-21 = 21-item Depression Anxiety Stress Scales; SD = standard deviation.