

Table 2 Incidence of metabolic syndrome in men and women, Zahedan, 2017

Population	Crude incidence rate, % (95% CI)				Age-standardized rate, % (95% CI)			
	AHA/NHLBI criteria	NCEP-ATP III criteria	ATP III criteria	IDF criteria	AHA/NHLBI criteria	NCEP-ATP III criteria	ATP III criteria	IDF criteria
Men	20.74 (16.25-25.83)	12.81 (9.35-16.97)	10.49 (7.37-14.35)	17.00 (12.92-21.73)	19.76 (15.33-24.74)	11.48 (8.27-15.58)	9.22 (6.33-12.95)	15.84 (12.03-20.64)
Women	35.27 (29.44-41.43)	29.56 (24.22-35.34)	27.50 (22.35-33.12)	35.00 (29.07-41.04)	34.80 (29.07-41.04)	28.70 (23.54-34.58)	26.86 (21.69-32.37)	34.70 (29.07-41.04)
Total	25.73 (23.84-31.46)	20.53 (17.35-24.01)	18.37 (15.36-21.70)	25.26 (21.71-29.08)	27.18 (23.50-31.09)	19.73 (16.57-23.12)	17.21 (14.29-20.47)	24.48 (21.03-28.34)

CI = confidence interval; AHA/NHLBI = American Heart Association and National Heart, Lung and Blood Institute; NCEP-ATP = National Cholesterol Education Program-Adult Treatment Panel; IDF = International Diabetes Federation.