

Table 1 Percent change in the utilization of key RMNCH services during various phases of COVID-19 pandemic in 2020

| RMNCH indicators | RMNCH service users | | | | | Percent change | | |
|--|---------------------|----------|----------|----------|--------------|----------------|--------------|--|
| | Jan 2020 | May 2020 | Sep 2020 | Dec 2020 | Jan–May 2020 | June–Sep 2020 | Oct–Dec 2020 | |
| Pre-pregnancy | | | | | | | | |
| Family planning visits | 283 760 | 141 057 | 191 314 | 163 751 | -50.3% | 17.7% | -9.7% | |
| Pregnancy | | | | | | | | |
| ANC-1 | 505 179 | 325 733 | 539 716 | 389 620 | -35.5% | 42.4% | -29.7% | |
| TT2 vaccine | 284 943 | 233 667 | 309 743 | 244 316 | -18.0% | 26.7% | -23.0% | |
| Delivery | | | | | | | | |
| Institutional Deliveries | 172 588 | 108 100 | 74 875 | 61 350 | -37.4% | -19.3% | -7.8% | |
| C section | 24 778 | 10 761 | 23 692 | 16 360 | -56.6% | 52.2% | -29.6% | |
| Postnatal | | | | | | | | |
| PNC-1 | 204 033 | 127 460 | 218 757 | 189 889 | -37.5% | 44.7% | -14.1% | |
| Infancy and childhood | | | | | | | | |
| Children < 12 mo received 3rd pentavalent vaccinea | 365 697 | 231 859 | 393 875 | 384 716 | -36.6% | 44.3% | -2.5% | |
| Reported cases of pneumonia < 5 yr | 154 415 | 42 450 | 97 428 | 95 413 | -72.5% | 35.6% | -1.3% | |
| Reported cases of diarrhoea < 5 yr | 428 022 | 381 061 | 653 436 | 353 729 | -11.0% | 63.6% | -70.0% | |
| Pneumonia < 5 yr inpatient treatment | 22 741 | 4202 | 6817 | 6251 | -81.5% | 11.5% | -2.5% | |

aVaccine used by the Expanded Program on Immunization in Pakistan to protect against diphtheria, pertussis, tetanus, hepatitis B and Haemophilus influenzae type B.

ANC-1 = first antenatal care visit; PNC-1 = first postnatal care visit; RMNCH = reproductive, maternal, newborn and child health; TT2 = second dose of tetanus toxoid.